

6450 U.S. Highway 1 Rockledge, FL 32955 myHFHP.org

# **Commercial Plans**

# 2022 Prior Authorization and Step Therapy Criteria

Health First Commercial Plans, Inc. is doing business under the name of Health First Health Plans. Health First Health Plans does not discriminate on the basis of race, color, national origin, disability, age, sex, gender identity, sexual orientation, or health status in the administration of the plan, including enrollment and benefit determinations.

36194\_MPINFO7674(10/19)

# Drugs

acitretin

#### **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

Severely impaired liver or kidney function. Chronic abnormally elevated blood lipid values. Concomitant use of methotrexate or tetracyclines. Pregnancy. Females of child-bearing potential who intend to become pregnant during therapy or at any time for at least 3 years after discontinuing therapy. Females of child-bearing potential who will not use reliable contraception while undergoing treatment and for at least 3 years following discontinuation. Females of child-bearing potential who drink alcohol during treatment or for two months after cessation of therapy.

#### **Required Medical Information**

Documented diagnosis of severe psoriasis

Age Restriction 18 years of age or older

**Prescriber Restriction** 

**Coverage Duration** Plan year

#### Other Criteria

Will not be approved for the treatment of acne.

# **ACTIMMUNE (interferon gamma-1B)**

#### Drugs ACTIMMUNE

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

Hypersensitivity to interferon gamma, E. coli derived proteins, or any component of the formulation.

# **Required Medical Information**

Documentation of diagnosis of chronic granulomatous disease or severe malignant osteoporosis.

Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Plan year

# ADCIRCA (tadalafil (Pulmonary Hypertension))

#### Drugs

tadalafil (pulm. hypertension)

**Covered Uses** 

All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

Receiving nitrate therapy (includes intermittent use).

# **Required Medical Information**

Age Restriction 18 years or older

**Prescriber Restriction** Cardiologist or Pulmonologist

**Coverage Duration** Plan year

# **Other Criteria**

Sildenafil citrate (generic Revatio indicated for Pulmonary Hypertension) must be tried first.

**Drugs** adefovir

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

**Required Medical Information** Clinical documentation of FDA approved indication for treatment.

Age Restriction 12 years and older.

**Prescriber Restriction** 

**Coverage Duration** Plan year

**Other Criteria** Optimal treatment duration is unknown.

#### Drugs ADEMPAS

#### **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

Pregnancy. Concomitant administration with nitrates or nitric oxide donors (such as amyl nitrate) in any form. Concomitant administration with phosphodiesterase inhibitors, including specific PDE-5 inhibitors (such as sildenafil, tadalafil, or vardenafil) or non-specific PDE inhibitors (such as dipyridamole or theophylline).

#### **Required Medical Information**

Age Restriction 18 years of age or older

**Prescriber Restriction** 

**Coverage Duration** Plan year

#### **Other Criteria**

For renewal, medication was effective (i.e. improved 6 minute walk distance, oxygen saturation, etc.)

Drugs AFINITOR ORAL TABLET 10 MG, 2.5 MG, 5 MG, 7.5 MG, everolimus (antineoplastic) oral tablet 2.5 mg, 5 mg, 7.5 mg

# **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

# **Required Medical Information**

Diagnosis of one of the following: A) Advanced metastatic renal cell carcinoma and patient has failed therapy (disease progressed) with Sutent or Nexavar (Afinitor only), or B) Progressive pancreatic, nonfunctional GI or lung neuroendocrine tumors (NET) that are unresectable, locally advanced or metastatic (Afinitor only), or C) Renal angiomyolipoma with tuberous sclerosis complex (TSC) and patient does not require immediate surgery (Afinitor only), or D) Advanced hormone receptor-positive, HER2-negative breast cancer and patient is a postmenopausal woman and patient has failed treatment with Femara or Arimidex and the medication will be used in combination with Aromasin (Afinitor only), or E) Subependymal giant cell astrocytoma (SEGA) associated with TSC that requires therapeutic intervention but is not a candidate for curative surgical resection (Afinitor or Afinitor Disperz only).

# **Age Restriction**

18 years of age or older for RCC, pNET, and renal angiomyolipoma with TSC. 1 year of age or older for SEGA.

**Prescriber Restriction** 

Coverage Duration Plan year

Drugs ALECENSA

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Documented diagnosis of anaplastic lymphoma kinase (ALK)-positive metastatic non-small cell lung cancer (NSCLC) as detected by an FDA-approved test.

# Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Through end of benefit year

Drugs ALINIA

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

# **Required Medical Information**

Clinical documentation indicating treatment needed for diarrhea caused by Giardia lamblia or Cryptosporidium parvum.

Age Restriction Age 1 year or older (Suspension) Age 12 years or older (Tablets)

**Prescriber Restriction** 

**Coverage Duration** 30 days

# Drugs ALIQOPA

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

**Required Medical Information** Documented diagnosis of relapsed follicular lymphoma (FL) who have received at least two prior systemic therapies.

# Age Restriction

**Prescriber Restriction** Oncologist or Hematologist

**Coverage Duration** Through end of benefit year

#### Drugs

alosetron

# **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

Initial therapy for Irritable Bowel Syndrome (IBS) in the male gender.

# **Required Medical Information**

Initial Therapy for Irritable Bowel Syndrome (IBS): 1. Confirmed diagnosis of IBS with diarrhea predominant symptoms for at least 6 months Reauthorization for Irritable Bowel Syndrome (IBS): 1. Recurrence of diarrhea predominant IBS, AND 2. documentation of positive clinical response while on Lotronex.

# **Age Restriction**

Patient must be at least 18 years of age or older

# **Prescriber Restriction**

Prescriber must be specially trained gastrointestinal physician

# **Coverage Duration**

IBS Initial Therapy: 12 weeks Reauthorization: 6 months

#### **Other Criteria**

Initial Therapy for Irritable Bowel Syndrome (IBS): 1. Failure to both: a. An antispasmodic (e.g. dicyclomine) AND b. An anti-diarrhea agent (e.g. loperamide, diphenoxylate-atropine)

# Alunbrig

#### Drugs ALUNBRIG ORAL TABLET 180 MG, 30 MG, 90 MG, ALUNBRIG ORAL TABLETS, DOSE PACK

# **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

diagnosis of A) anaplastic lymphoma kinase (ALK)-positive metastatic non-small cell lung cancer (NSCLC) who have progressed on or are intolerant to crizotinib OR B) tyrosine kinase inhibitor-naive advanced anaplastic lymphoma kinase (ALK)-positive metastatic non-small cell lung cancer (NSCLC).

# Age Restriction

Prescriber Restriction Oncologist

**Coverage Duration** 3 Months

# **AMPYRA** (dalfampridine)

#### Drugs

dalfampridine

# **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

History of seizure. Moderate or severe renal impairment (creatinine clearance less than or equal to 50 mL/minute). Patient currently using any other forms of 4-aminopyridine.

#### **Required Medical Information**

Diagnosis of multiple sclerosis AND patient is ambulatory AND patient has walking impairment.

#### **Age Restriction**

Prescriber Restriction Neurologist

# Coverage Duration

Initial: 3 months. Renewal: Plan year

#### **Other Criteria**

For renewal, documentation that walking speed has improved from baseline must be provided.

Drugs APTIOM

**Covered Uses** All FDA-approved indications not otherwise excluded from Health Plan.

**Exclusion Criteria** 

**Required Medical Information** 

Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Plan year

#### Drugs APTIVUS

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

# **Required Medical Information** Clinical documentation of FDA approved indication for treatment.

Age Restriction 2 years of age and older

**Prescriber Restriction** 

**Coverage Duration** Plan year

**Other Criteria** Do not use Aptivus in treatment-naïve patients.

#### Drugs ARCALYST

# **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

Use in combination with other IL-1 inhibitors (e.g. Ilaris, Kineret) or tumor necrosis factor (TNF) inhibitors (e.g. Enbrel, Humira, Remicade, etc). Individual is receiving live vaccines. Exhibiting evidence of active or chronic infection(s), including tuberculosis, or a history of recurrent infections. Individual has not had a tuberculin skin test (TST) or Centers for Disease Control (CDC)-recommended equivalent to evaluate for latent tuberculosis prior to initiating treatmentwith rilonacept.

# **Required Medical Information**

Documented diagnosis of cryopyrin-associated period syndromes (CAPS), including Familial Cold Auto-inflammatory Syndrome (FCAS) and/or Muckle-Wells Syndrome (MWS). Member's diagnosis of CAPS must be confirmed by either NRLP=3 gene mutation OR overproduction of interleukin-1.

# Age Restriction

12 years of age and older

# **Prescriber Restriction**

Prescribed by or in consultation with or recommendation of, an immunologist, allergist, dermatologist, rheumatologist, neurologist, or other medical specialist.

**Coverage Duration** Plan year

Other Criteria Approve doses based on FDA labeling.

# Armodafinil (NUVIGIL)

#### Drugs

armodafinil

#### **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

# Required Medical Information

Documentation of diagnosis and treatment history.

Age Restriction Patient must be at least 17 years or older

# **Prescriber Restriction**

Idiopathic hypersomnia-- approve if the diagnosis is confirmed by a sleep specialist physician or at an institution that specializes in sleep disorders (i.e., sleep center)

**Coverage Duration** Plan year

#### Drugs ATRIPLA

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

# Required Medical Information

Clinical documentation of FDA approved indication for treatment.

Age Restriction 12 years and older

# **Prescriber Restriction**

**Coverage Duration** Plan year

Drugs AUBAGIO

**Covered Uses** All FDA-approved indications not otherwise excluded from Health Plan

# **Exclusion Criteria**

# **Required Medical Information**

Diagnosis of relapsing forms of multiple sclerosis and must have tried and failed one generic alternative agent such as dimethyl fumarate or glatiramer acetate.

# Age Restriction

**Prescriber Restriction** Prescribed by or in consultation with a neurologist or MS specialist.

#### **Coverage Duration** Plan year

# Drugs AUSTEDO ORAL TABLET 12 MG, 6 MG, 9 MG

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

**Exclusion Criteria** 

**Required Medical Information** 

Age Restriction

**Prescriber Restriction** 

**Coverage Duration** 3 Months

#### Drugs

# AVONEX INTRAMUSCULAR PEN INJECTOR, AVONEX INTRAMUSCULAR PEN INJECTOR KIT, AVONEX INTRAMUSCULAR SYRINGE, AVONEX INTRAMUSCULAR SYRINGE KIT

#### **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

Treatment of primary progressive MS is not covered. Combination therapy with a beta interferon product, Gilenya, Aubagio, Tecfidera, Tysabri or Copaxone is not covered.

#### **Required Medical Information**

Diagnosis of relapsing forms of multiple sclerosis and must have tried and failed one generic alternative agent such as dimethyl fumarate or glatiramer acetate

#### **Age Restriction**

#### **Prescriber Restriction**

Prescribed by or in consultation with a neurologist or MS specialist.

**Coverage Duration** Plan year

Drugs AYVAKIT ORAL TABLET 100 MG, 200 MG, 300 MG

**Covered Uses** 

**Exclusion Criteria** 

# **Required Medical Information**

documentation of unresectable or metastatic GIST harboring a platelet-derived growth factor receptor alpha (PDGFRA) exon 18 mutation.

Age Restriction

Prescriber Restriction Oncologist, Allergist, Immunologist

**Coverage Duration** 3 months

Drugs BALVERSA

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Documented diagnosis of locally advanced or metastatic urothelial carcinoma AND member has susceptible FGFR3 or FGFR2 genetic alteration as detected by an FDA-approved companion diagnostic AND disease has progressed during or following at least one line of prior platinum-containing chemotherapy including within 12 months of neoadjuvant or adjuvant platinum-containing chemotherapy.

Age Restriction 8 years or older

**Prescriber Restriction** 

**Coverage Duration** 3 Months

#### Drugs BANZEL ORAL SUSPENSION, BANZEL ORAL TABLET 200 MG, 400 MG, rufinamide oral suspension

# **Covered Uses**

All FDA-approved indications not otherwise excluded from Health Plan.

#### **Exclusion Criteria**

Banzel is not covered for members with the diagnosis of Familial Short QT syndrome

#### **Required Medical Information**

Documentation of diagnosis. Documentation of previous therapies and that the current medication regimen is inadequate to control disease.

# Age Restriction

Must be 1 years of age or older

# **Prescriber Restriction**

**Coverage Duration** Plan year

#### **Other Criteria**

Patient must be refractory to at least 2 of the following: Felbamate (Felbatol), Lamotrigine (Lamictal), Topiramate (Topamax), Valproic acid (Depakene), Divalproex sodium (Depakote)

# Drugs BARACLUDE ORAL SOLUTION, entecavir

**Covered Uses** 

All FDA-approved indications not otherwise excluded by Health Plan.

**Exclusion Criteria** 

**Required Medical Information** Documentation of diagnosis

**Age Restriction** 

**Prescriber Restriction** 

Coverage Duration Plan year

# Drugs BAVENCIO

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

# Age Restriction

Prescriber Restriction Oncologist

**Coverage Duration** 3 Months

# Drugs BELSOMRA

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

Required Medical Information Documented trial and failure of two formulary alternatives AND documented medication review to rule out medication induced insomnia.

# **Age Restriction**

**Prescriber Restriction** 

**Coverage Duration** Through end of benefit year

# BENLYSTA

#### Drugs BENLYSTA SUBCUTANEOUS AUTO-INJECTOR

**Covered Uses** All FDA-approved indications not otherwise exluded for part D

**Exclusion Criteria** 

**Required Medical Information** Documentation from the medical record of diagnosis

Age Restriction

**Prescriber Restriction** 

**Coverage Duration** 3 months

# Benznidazole

Drugs

benznidazole

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

#### **Exclusion Criteria**

Patients who have used disulfiram within two weeks of initiation of benznidazole

#### **Required Medical Information**

Documentation of a consultation with an infectious disease specialist. Reviewer will verify patient claim history to confirm that patient has not used disulfiram within two weeks prior to benznidazole initiation

**Age Restriction** 

**Prescriber Restriction** 

**Coverage Duration** 60 days

# **Benzodiazepines**

# Drugs DIAZEPAM INTENSOL, diazepam oral concentrate

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

**Required Medical Information** Documentation of diagnosis

Age Restriction

**Prescriber Restriction** 

Coverage Duration Plan year

#### Drugs BETASERON SUBCUTANEOUS KIT

# **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

Treatment of primary progressive MS is not covered. Combination therapy with a beta interferon product, Gilenya, Aubagio, Tecfidera, Tysabri or Copaxone is not covered.

# **Required Medical Information**

Diagnosis of relapsing form of multiple sclerosis OR diagnosis of first clinical episode and MRI features consistent with multiple sclerosis.

# Age Restriction

#### **Prescriber Restriction**

Prescribed by or in consultation with a neurologist or MS specialist.

# **Coverage Duration**

Plan year

# Other Criteria

Approve if: 1) Patient is currently taking Betaseron, OR 2) Patient has tried dimethyl fumarate (Tecfidera), interferon beta-1a intramuscular (Avonex) and glatiramer acetate (Copaxone)

Drugs BLENREP

Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

1. Member has been diagnosed with relapsed or refractory multiple myeloma AND 2. Member had disease progression on at least four prior anti-myeloma treatment (including a proteasome inhibitor, immunomodulatory agent, and anti-CD38 monoclonal antibody AND 3. Member has an ophthalmic exam (i.e. visual acuity) at baseline, prior to each infusion AND 4 Member has no current corneal epithelial disease (excluding mild punctatekeratopathy) AND 5 Member has not had a prior allogeneic stem cell transplant AND 6. Member does not have active POEMS syndrome or active plasma cell leukemia AND 7. Member enrolled in BLENREP REM program AND 8. Therapy will be used in combination with preservative-free lubricant eye drops AND 9. medication will be used as single-agent therapy

# **Required Medical Information**

Age Restriction Member is age 18 years or older

Prescriber Restriction Oncologist

**Coverage Duration** 3 months

# BOSULIF

#### Drugs BOSULIF ORAL TABLET 100 MG, 400 MG, 500 MG

#### **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

Coverage is not provided in those that have the BCR-ABL1 T315I mutation and use of Bosulif in combination with other kinase inhibitors (for example sorafenib, sunitinib, etc.)

# **Required Medical Information**

Diagnosis of Philadelphia chromosome-positive chronic myelogenous leukemia.

Age Restriction 18 years of age and older

Prescriber Restriction Hematologist/Oncologist

**Coverage Duration** 3 Months

#### Drugs BRAFTOVI ORAL CAPSULE 50 MG, 75 MG

# **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Documentation of unresectable or metastatic melanoma with a BRAF V600E or V600K mutation, as detected by an FDAapproved test and the medication will be used in combination with binimetinib.

# **Age Restriction**

Prescriber Restriction Oncology

**Coverage Duration** 3 months

Drugs CAPLYTA

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

**Required Medical Information** Documented trial of any of the two following drugs: aripiprazole, clozapine, olanzapine, risperidone, quetiapine, ziprasidone

**Age Restriction** 

**Prescriber Restriction** 

**Coverage Duration** Through end of benefit year

#### Drugs

raloxifene, tamoxifen

#### **Covered Uses**

This criteria is a copay review process. The medications tamoxifen or raloxifene may be eligible for \$0 cost-share for women 35 years of age or older who: 1) do not have a history of breast cancer, and 2) are being prescribed tamoxifen or raloxifene for the purpose of primary prevention of invasive breast cancer because the member is deemed high risk, and 3) are post-menopausal, if prescribed raloxifene (this requirement does not apply to tamoxifen)

#### **Exclusion Criteria**

Women under 35 years of age, history of breast cancer

#### **Required Medical Information**

A 5-year predicted risk of breast cancer greater than or equal to 1.66%, as calculated by the Gail model.

Age Restriction 35 years and older

**Prescriber Restriction** 

**Coverage Duration** 5 years

Drugs BROVANA

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

**Required Medical Information** Documentation of diagnosis

Age Restriction 18 years and older.

**Prescriber Restriction** 

Coverage Duration Plan year

# **Other Criteria**

Must have documented failure, intolerance or contraindication to a long-acting beta agonist formulary product OR be unable to use a hand-actuated device.

Drugs BRUKINSA

**Covered Uses** 

**Exclusion Criteria** 

#### **Required Medical Information**

Documentation of mantle cell lymphoma (MCL) in patients who have received at least one prior therapy.

Age Restriction

Prescriber Restriction Oncology

**Coverage Duration** 3 Months

Drugs BUPHENYL ORAL TABLET, sodium phenylbutyrate oral tablet

# **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

# **Required Medical Information**

Documentation of diagnosis confirmed by enzymatic, biochemical or genetic testing. Buphenyl will be used for chronic management of urea cycle disorders (UCD).

# Age Restriction

**Prescriber Restriction** 

Coverage Duration Plan year

#### **Buprenorphine Products**

#### Drugs

buprenorphine HCl sublingual, buprenorphine-naloxone sublingual film 12-3 mg, 2-0.5 mg, 4-1 mg, 8-2 mg

#### Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

#### **Exclusion Criteria**

Being used for the treatment of pain OR patient is using short or long acting narcotics concurrently with Suboxone/Subutex.

#### **Required Medical Information**

The indicated diagnosis and medication usage must be supported by documentation from the patient's medical records.

#### Age Restriction

Must be 16 years of age or older.

#### **Prescriber Restriction**

Prescribing provider must have a DEA number starting with the letter X, AND physician must be listed on the Buprenorphine Physician Locator maintained by the Substance Abuse and Mental Health Services Administration (SAMSHA).

**Coverage Duration** 12 months

Drugs BUSULFEX

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

**Exclusion Criteria** 

**Required Medical Information** Documentation of diagnosis

Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Plan year

# Drugs CABOMETYX ORAL TABLET 20 MG, 40 MG, 60 MG

#### **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

#### **Required Medical Information**

Diagnosis of advanced renal cell carcinoma (RCC) or hepatocellular carcinoma in patients who have been previously treated with sorafenib.

#### Age Restriction

**Prescriber Restriction** 

**Coverage Duration** 3 months

#### Drugs AIMOVIG AUTOINJECTOR, AJOVY AUTOINJECTOR, AJOVY SYRINGE

# **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Clinical documentation of migraines and member has tried and failed two formulary alternatives for migraine prophylaxis with two different mechanism of action.

# Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Through end of benefit year

#### Drugs CALQUENCE

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

**Required Medical Information** Documented diagnosis of mantle cell lymphoma (MCL) who have received at least one prior therapy.

# Age Restriction

**Prescriber Restriction** Oncologist or hematologist

**Coverage Duration** 3 Months

**Drugs** capecitabine

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

**Required Medical Information** Clinical documentation of FDA approved indication for treatment.

Age Restriction 18 years and older.

**Prescriber Restriction** Oncologist

**Coverage Duration** Plan year

# **CAPRELSA** (vandetanib)

#### Drugs CAPRELSA ORAL TABLET 100 MG, 300 MG

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria Congenital long QT syndrome

#### **Required Medical Information**

Diagnosis of symptomatic or progressive medullary thyroid cancer with unresectable locally advanced or metastatic disease.

Age Restriction 18 years or older

**Prescriber Restriction** Oncologist or endocrinologist

**Coverage Duration** 3 Months

#### CASPOFUNGIN

Drugs

caspofungin

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

#### **Exclusion Criteria**

#### **Required Medical Information**

Diagnosis of one of the following: A)Empirical therapy for presumed fungal infections in febrile, neutropenic patients, or B)Treatment of candidemia and other Candida infections (intraabdominal abscesses, peritonitis and pleural space infections), or C)Treatment of esophageal candidiasis, or D)Treatment of invasive aspergillosis in patients who are refractory to or intolerant of other therapies (amphotericin B, itraconazole).Age Restrictions: 3 months of age or older

#### **Age Restriction**

**Prescriber Restriction** 

**Coverage Duration** 3 months

#### Drugs CAYSTON

# **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D. All FDA-approved indications not otherwise excluded from Health Plan.

# **Exclusion Criteria**

#### **Required Medical Information**

Diagnosis of cystic fibrosis AND patient has evidence of P. aeruginosa in the lungs

Age Restriction 7 years of age and older

#### **Prescriber Restriction**

**Coverage Duration** Plan year

# **Other Criteria**

For renewal, Patient is benefiting from treatment (i.e. improvement in lung function [FEV1], decreased number of pulmonary exacerbations)

cinacalcet

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Documentation of secondary hyperparathyroidism due to chronic kidney disease on dialysis, Or Hypercalcemia due to parathyroid carcinoma, Or severe hypercalcemia in patients with primary hyperparathyroidism who are unable to undergo parathyroidectomy.

**Age Restriction** 

**Prescriber Restriction** 

**Coverage Duration** 3 months

clozapine oral tablet, disintegrating

#### **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

If the patient has any of the following contraindications: agranulocytosis, bone marrow suppression, coma, ileus, leukopenia, myocarditis or neutropenia, OR if the patient has CNS depression, dementia-related psychosis or uncontrolled epilepsy.

#### **Required Medical Information**

A statement showing the patient is unwilling or unable to take tablets or capsules orally or at high risk for non-compliance AND is not receiving other tablets or capsules indicating that the patient can take non-dissolvable tablets.

**Age Restriction** 

**Prescriber Restriction** Part of a clozapine registry.

**Coverage Duration** Plan Year

#### COMETRIQ

```
Drugs
COMETRIQ ORAL CAPSULE 100 MG/DAY(80 MG X1-20 MG X1), 140 MG/DAY(80 MG X1-20 MG X3), 60 MG/DAY
(20 MG X 3/DAY)
```

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

**Exclusion Criteria** Gastrointestinal perforation. Fistula. Severe hemorrhage.

# Required Medical Information

Diagnosis of progressive metastatic, medullary thyroid cancer.

Age Restriction 18 years or older

Prescriber Restriction Oncologist/Hematologist

**Coverage Duration** 3 Months

#### Drugs COPAXONE SUBCUTANEOUS SYRINGE 20 MG/ML, 40 MG/ML

#### **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

Treatment of primary progressive MS is not covered. Combination therapy with a beta interferon product, Gilenya, Aubagio, Tecfidera, Tysabri or Copaxone is not covered.

#### **Required Medical Information**

Diagnosis of relapsing forms of multiple sclerosis and must have tried and failed one generic alternative agent such as dimethyl fumarate or glatiramer acetate

#### Age Restriction

**Prescriber Restriction** By or in consultation with a Neurologist or a Certified MS Specialist

**Coverage Duration** Plan year

Drugs COPIKTRA

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Documentation of the following: A) relapsed or refractory CLL or SLL after at least two prior therapies OR B)relapsed or refractory FL after at least two prior systemic therapies.

# Age Restriction

Prescriber Restriction Oncology

**Coverage Duration** 3 months

#### Drugs CORLANOR ORAL SOLUTION

# **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Must be clinically diagnosed with A) stable, symptomatic chronic heart failure in adults with left ventricular ejection fraction less than or equal to 35% supported by documentation from the patient's medical records, AND have sinus rhythm with resting heart rate greater than or equal to 70 beats per minute, AND be on maximally tolerated doses of beta blockers unless contraindicated, AND be on optimal therapy with standard treatment of ACEI or ARB unless intolerant or contraindicated, AND be on optimal therapy with standard treatment of an aldosterone antagonist unless intolerant or contraindicated, AND documentation of trial and failure of Entresto OR B)stable symptomatic heart failure due to dilated cardiomyopathy (DCM) in pediatric patients aged 6 months and older AND have sinus rhythm with an elevated heart rate.

# Age Restriction

Prescriber Restriction Cardiologist

**Coverage Duration** Through end of benefit year

#### Drugs COSENTYX, COSENTYX (2 SYRINGES), COSENTYX PEN, COSENTYX PEN (2 PENS)

# **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Diagnosis of 1 of the following: A) Mod-severe chronic plaque psoriasis (affecting more than 5% of body surface area or crucial body areas such as the hands, feet, face, or genitals) and trial of at least 1 of the following: phototherapy (including but not limited to Ultraviolet A with a psoralen [PUVA] and/or retinoids [RePUVA] for at least 1 continuous month or 1 or more oral systemic treatments (e.g., MTX, cyclosporine, acitretin, sulfasalazine) for at least 3 months, OR B) Psoriatic arthritis and patient had an inadequate response, intolerance, or contraindication to methotrexate, OR C) Ankylosing spondylitis and patient had an inadequate response, intolerance or contraindication to at least two NSAIDs, OR D) Non-radiographic Axial Spondyloarthritis and patient had an inadequate response, intolerance or contraindication to at least two NSAIDs.

Age Restriction

**Prescriber Restriction** 

**Coverage Duration** 12 months.

Drugs COTELLIC

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Documented unresectable or metastatic melanoma with a BRAF V600E or V600K mutation, in combination with vemurafenib.

Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Through end of benefit year.

Drugs CYCLOSET

**Covered Uses** All FDA-approved indications not otherwise excluded from Health Plan

**Exclusion Criteria** 

**Required Medical Information** Documented diagnosis of type 2 diabetes mellitus

Age Restriction

**Prescriber Restriction** 

Coverage Duration Plan year

#### Drugs CYSTAGON

#### **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

Cysteamine is contraindicated in patients who have demonstrated hypersensitivity to cysteamine or penicillamine hypersensitivity.

# **Required Medical Information**

Documentation of diagnosis

#### Age Restriction

#### **Prescriber Restriction**

# **Coverage Duration**

3 Months

# **Other Criteria**

Do not administer intact cysteamine capsules to children less than 6 years old because of aspiration risk. Capsules may be administered by sprinkling contents over food.

Drugs DANYELZA

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

1. Medication will be used in combination with granulocyte-macrophage colony-stimulating factor (GM-CSF) AND 2 documented diagnosis of pediatric patients 1 year of age and older and adult patients with relapsed or refractory high-risk neuroblastoma in the bone or bone marrow AND 3. patient has demonstrated a partial response, minor response, or stable disease to prior therapy.

# Age Restriction

Prescriber Restriction Oncologist

**Coverage Duration** 3 months

#### Drugs DAURISMO ORAL TABLET 100 MG, 25 MG

#### **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D.

#### **Exclusion Criteria**

#### **Required Medical Information**

Documentation that the medication will be used in combination with low-dose cytarabine for the treatment of newlydiagnosed acute myeloid leukemia (AML), and the patient is A) 75 years of age or older old OR B)has comorbidities that preclude use of intensive induction chemotherapy.

#### Age Restriction

Prescriber Restriction oncology

**Coverage Duration** 3 months

Drugs DESCOVY ORAL TABLET 200-25 MG

**Covered Uses** All FDA-approved indications not otherwise excluded from Health Plan

**Exclusion Criteria** 

**Required Medical Information** 

Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Plan year

# Drugs DIACOMIT ORAL CAPSULE

# **Covered Uses**

# **Exclusion Criteria**

# **Required Medical Information**

Documented diagnosis of Dravet syndrome. Not appropriate for monotherapy. DIACOMIT is to be used as an add-on therapy to clobazam for seizures associated with Dravet syndrome when these are not adequately controlled.

# Age Restriction 2 years and older

# **Prescriber Restriction**

**Coverage Duration** Through end of benefit year

# **Other Criteria**

Dosing should not exceed 50 mg/kg/day, administered by mouth in 2or 3 divided doses.

diclofenac sodium topical gel 3 %

# **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

**Required Medical Information** Documentation of diagnosis and treatment history

# Age Restriction DO NOT use Solaraze in children.

# **Prescriber Restriction**

**Coverage Duration** 3 months

**Other Criteria** Must have failed topical 5-FU cream

#### **Didanosine (VIDEX)**

#### Drugs

didanosine oral capsule, delayed release (DR/EC) 250 mg, 400 mg

# **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

#### **Required Medical Information**

Clinical documentation of FDA approved indication for treatment.

Age Restriction 2 weeks and older.

# **Prescriber Restriction**

**Coverage Duration** Plan year

# Drugs DIFICID ORAL TABLET

# **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

# **Required Medical Information**

Diagnosis of Clostridium difficile associated diarrhea (CDAD) with one of the following: A) Patient has mild to moderate CDAD and failure, contraindication or intolerance to oral Flagyl (metronidazole) and oral Vancocin (vancomycin), or B) Patient has severe CDAD.

Age Restriction 18 years or older

**Prescriber Restriction** 

**Coverage Duration** 10 Days

doxercalciferol oral

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

# **Required Medical Information** Clinical documentation of FDA approved indication for treatment.

Age Destriction

Age Restriction 18 years and older.

**Prescriber Restriction** 

**Coverage Duration** Plan year

dronabinol oral capsule 10 mg, 2.5 mg, 5 mg

#### **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

Any pertinent clinical situation as defined by the product label that could affect patient safety and/or therapeutic efficacy (i.e. contraindications, warnings, precautions, adverse effects, renal or hepatic function, drug interactions, lab values, required prior or concomitant therapy, inappropriate dosing and/or duration, etc).

#### **Required Medical Information**

For treatment of chemotherapy-induced nausea or vomiting refractory to conventional antiemetic agents: 1. Patient is receiving cancer chemotherapy, AND 2. Failure to preferred 5HT-3 receptor antagonist. preferred agents include ondansetron or granisetron, AND 3. Failure to one of the following agents: a. Antihistamine b. Corticosteroid c. Prokinetic agent d. Antipsychotic. For treatment of anorexia associated with weight loss in patients with HIV: documentation of trial and failure, contraindication, or intolerance to megestrol.

#### **Age Restriction**

18 years old and greater for the treatment of anorexia associated with weight loss in patients with HIV

**Prescriber Restriction** 

**Coverage Duration** 3 Months

# DUPIXENT PEN SUBCUTANEOUS PEN INJECTOR 300 MG/2 ML, DUPIXENT SYRINGE SUBCUTANEOUS SYRINGE 200 MG/1.14 ML, 300 MG/2 ML

# **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Patient must have the following: A) Moderate-to-severe atopic dermatitis (eczema)AND submission of medical records (e.g. chart notes, laboratory values) documenting the following: Inadequate response, intolerance or contraindication to ONE medication in EACH of the following categories:a. Topical calcineurin inhibitor b. High potency topical corticosteroid. OR B) Moderate-to-severe asthma AND submission of medical records documenting the following: 1. Patient has ONE of the following: a. Asthma with eosinophilic phenotype with eosinophil count greater than or equal to 300 cells/mcL in the past 12 months, or b. Oral corticosteroid dependent asthma with at least 1 month of daily oral corticosteroid use within the last 3 months AND 2. Inadequate control of asthma symptoms after a minimum of 3 months of compliant use of one of the following: a. Inhaled corticosteroids & long acting beta2 agonist, or b. Inhaled corticosteroids & long acting muscarinic antagonist. OR C)Chronic rhinosinusitis with nasal polyposis (CRSwNP)AND submission of medical records (e.g. chart notes, laboratory values) documenting the following: Inadequate response, intolerance or contraindication to ONE medication in EACH of the following categories: 1. Inadequate response, intolerance or contraindication to ONE medication in EACH of the following categories: a. Nasal corticosteroid spray and b. Oral corticosteroid.

# Age Restriction

18 or older

#### Prescriber Restriction

Dermatologist or allergist/immunologist

#### Coverage Duration

Initial Authorization will be for 3 months. Reauthorization will be for 1 year.

#### **Other Criteria**

Renewals require submission of medical records (e.g. chart notes, laboratory values) documenting improvement of the condition.

# **EDECRIN** (Ethacrynic Acid)

#### Drugs EDECRIN

**Covered Uses** All FDA-approved indications not otherwise excluded from Health Plan

**Exclusion Criteria** Use for pediatrics 12 months and younger

# **Required Medical Information**

Documentation of diagnosis.

Age Restriction 13 months and older.

#### **Prescriber Restriction**

**Coverage Duration** Plan year

#### **Other Criteria**

Must have documented failure, intolerance or contraindication to at least 2 other loop diuretics.

# Drugs EDURANT

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

**Required Medical Information** Clinical documentation of FDA approved indication for treatment.

# Age Restriction 12 years and older.

# **Prescriber Restriction**

**Coverage Duration** Plan year

Drugs ELZONRIS

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Patient must have a definitive diagnosis of Blastic Plasmacytoid Dendritic Cell Neoplasm (BPDCN) AND Patient has CD123 positive expressing disease AND Patient has a baseline serum albumin level of at least 3.2 g per dL

Age Restriction Patient is 2 years of age or older

**Prescriber Restriction** 

**Coverage Duration** 6 months

#### Drugs EMCYT

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

#### **Required Medical Information** Documented diagnosis of metastatic and/or progressive prostate cancer.

Age Restriction 18 years or older

Prescriber Restriction Oncologist prescriber

**Coverage Duration** 6 months

#### Drugs EMEND ORAL CAPSULE 80 MG, EMEND ORAL CAPSULE, DOSE PACK

#### **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

# **Required Medical Information**

Clinical documentation of FDA approved indication for treatment.

# Age Restriction 18 years and older.

# **Prescriber Restriction**

**Coverage Duration** 3 Months

#### Drugs EMSAM

# **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

Pheochromocytoma. Patient is taking or will take any of the following: SSRIs, SNRIs, tricyclic antidepressants (TCAs), bupropion, buspirone, meperidine, tramadol, methadone, pentazocine, dextromethorphan, St. John's wort, mirtazapine, cyclobenzaprine, oral selegiline, other MAOIs, oxcarbazepine, carbamazepine, and/or sympathomimetic amines.

### **Required Medical Information**

Diagnosis of major depressive disorder, AND 1) Failure of at least two generic oral antidepressants from different classes(at least one should be from the following list: selective serotonin reuptake inhibitors, serotonin and norepinephrine reuptake inhibitors, mirtazapine, or bupropion unless contraindicated), AND 2) Patient had an adequate washout period (for patients previously on agents requiring a washout period)

Age Restriction

**Prescriber Restriction** 

Coverage Duration Plan year

Drugs EMTRIVA

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

**Exclusion Criteria** 

**Required Medical Information** Clinical documentation of FDA approved indication for treatment

Age Restriction

**Prescriber Restriction** 

Coverage Duration Plan year

# Drugs

# ENBREL SUBCUTANEOUS RECON SOLN, ENBREL SUBCUTANEOUS SOLUTION, ENBREL SUBCUTANEOUS SYRINGE 25 MG/0.5 ML (0.5), 50 MG/ML (1 ML), ENBREL SURECLICK

### **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

### **Exclusion Criteria**

Using in combination with other TNF antagonists, IL-1R antagonists, janus kinase inhibitor, anti-cd20 monoclonalantibodies or selective co-stimulation modulators.

# **Required Medical Information**

Diagnosis of one of the following: A) Moderate to severe rheumatoid arthritis and patient had an inadequate response to, intolerance to, or contraindication to at least one non-biologic disease modifying anti-rheumatic drugs (DMARD) and one NSAID for at least 3 months, or B) Moderate to severe polyarticular juvenile idiopathic arthritis and patient had an inadequate response, intolerance or contraindication to at least one DMARD and one NSAID for at least 3 months, OR C) Psoriatic arthritis and patient had an inadequate response, intolerance, or contraindication to methotrexate, or D) Ankylosing spondylitis and patient had an inadequate response, intolerance or contraindication to at least two NSAIDs, or E) Moderate to severe chronic plaque psoriasis (affecting more than 5% of body surface area or affecting crucial body areas such as the hands, feet, face, or genitals) and patient had an inadequate response, intolerance or contraindication to conventional therapy with to at least two of the following: phototherapy (including but not limited to Ultraviolet A with a psoralen [PUVA] and/or retinoids [RePUVA] for at least one continuous month or one or more oral systemic treatments (i.e. methotrexate, cyclosporine, acitretin, sulfasalazine) for at least 3 months.

# **Age Restriction**

2 years of age or older for JIA. 18 years of age or older for all other indications.

**Prescriber Restriction** 

**Coverage Duration** 12 months

Drugs ENDARI

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

**Exclusion Criteria** 

**Required Medical Information** Documented diagnosis of sickle cell disease.

Age Restriction 5 years and older

**Prescriber Restriction** 

**Coverage Duration** Through end of benefit year

Drugs ENSPRYNG

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

a. Diagnosis of neuromyelitis optica spectrum disorder (NMOSD)AND b. Patient has a positive serologic test for antiaquaporin-4 (AQP4) antibodies AND c. History of failure, contraindication, or intolerance to rituximab therapy AND d. One of the following:(1) History of one or more relapses that required rescue therapy during the previous 12 months OR (2) History of two or more relapses that required rescue therapy during the previous 24 months

# Age Restriction

# **Prescriber Restriction**

Prescribed by, or in consultation with, a neurologist

**Coverage Duration** 6 months

#### Drugs EPCLUSA ORAL TABLET 400-100 MG

# **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

Current alcohol, drug use or reinfection after 3 months of successful treatment. Alcohol urine metabolite and drug screen required.

# **Required Medical Information**

Provider must submit medical records documenting the diagnosis of chronic hepatitis C with genotype and subtype, if applicable (i.e., genotypes 1, 2, 3, 4, 5, or 6) AND submit medical records documenting viral load taken within 6 months of beginning therapy, AND submit medical records documenting F2-F4 fibrosis with a fibrosis score of 0.48 and up or be a documented health Care worker in direct patient care setting.

# Age Restriction

18 years of age and older

# **Prescriber Restriction**

Prescribed by, or in consultation with, a gastroenterologist, hepatologist, or infectious disease physician.

# **Coverage Duration**

12 weeks or as defined by current AASLD/IDSA guidance.

# **Other Criteria**

Mavyret, generic Ledipasvir/Sofosbuvir, or generic Sofosbuvir/Velpatasvir must be tried first in patients with chronic hepatitis C. Criteria and coverage durations will be applied consistent with current AASLD/IDSA guidance. Must have contraindication to or be unable to tolerate Mavyret, generic Ledipasvir/Sofosbuvir, and generic Sofosbuvir/Velpatasvir.

Drugs EPIDIOLEX

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Documentation of the following A) diagnosis of seizures associated with Lennox-Gastaut syndrome (LGS)or seizures associated with Dravet syndrome (DS): AND B) normal serum transaminases (ALT and AST) and total bilirubin levels: AND C) inadequate treatment response, intolerance, or contraindication to TWO generic antiepileptic medications (i.e. clobazam, Valproic acid, Lamotrigine, Levetiracetam, Topiramate, etc.)

# Age Restriction

2 years of age or older

**Prescriber Restriction** 

# Coverage Duration

Through end of benefit year

# **Other Criteria**

Dosing is within the FDA labeled dose of up to 20mg/kg/day.

# Drugs EPIVIR HBV ORAL SOLUTION

# **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

**Required Medical Information** Clinical documentation of FDA approved indication for treatment.

Age Restriction 3 months of age or older

# **Prescriber Restriction**

**Coverage Duration** Plan year

# **EPZICOM** (Abacavir/Lamivudine)

#### Drugs

abacavir-lamivudine, EPZICOM

#### Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

**Required Medical Information** Clinical documentation of FDA approved indication for treatment.

# Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Plan year

Other Criteria Documented negative HLA-B\*5701 screening.

#### Drugs ERIVEDGE

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Diagnosis of metastatic basal cell carcinoma OR Diagnosis of locally advanced basal cell carcinoma that has recurred following surgery or when the patient is not a candidate for surgery and radiation.

Age Restriction 18 years or older

**Prescriber Restriction** 

**Coverage Duration** 3 months

# Drugs ERLEADA

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# Required Medical Information

Documented diagnosis of non-metastatic, castration-resistant prostate cancer (NM-CRPC).

# Age Restriction

**Prescriber Restriction** 

**Coverage Duration** 3 Months

# Drugs ERTACZO

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

#### **Required Medical Information** Clinical documentation of FDA approved indication for treatment.

Age Restriction 12 years and older.

**Prescriber Restriction** 

**Coverage Duration** 4 weeks

**Other Criteria** Failure to generic topical antifungal medications.

#### Drugs ESBRIET

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

The patient has a diagnosis of idiopathic pulmonary fibrosis confirmed by a high resolution CT scan or biopsy AND the patient does not have evidence or suspicion of an alternative interstitial lung disease diagnosis AND liver function tests have been performed prior to start of therapy and base line PFTs provided.

# Age Restriction

# **Prescriber Restriction**

Prescribed by or in consultation with a pulmonologist

# **Coverage Duration**

Through benefit year

# **Other Criteria**

For renewal, patient experienced stabilization from baseline or a less than 10 percent decline in force vital capacity AND the patient has not experienced AST or ALT elevations greater than 5 times the upper limit of normal or greater than 3 times the upper limit of normal with signs or symptoms of severe liver damage.

Drugs EVENITY

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Diagnosis of Osteoporosis in a postmenopausal female AND one or more of the following: 1) History of osteoporotic fracture, or 2) Documented trial and failure of bisphosphonate or 3) Documented contraindication or intolerance to bisphosphonate therapy. Patient has not received more than 1 year of therapy with Evenity.

**Age Restriction** 

**Prescriber Restriction** 

**Coverage Duration** 12 months

#### Drugs AKYNZEO (FOSNETUPITANT) INTRAVENOUS RECON SOLN, AKYNZEO (NETUPITANT), SANCUSO, VALCHLOR, VARUBI, XERMELO

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information** Documentation that the medication would be used for an FDA approved indication.

# Age Restriction

Prescriber Restriction Oncologist

**Coverage Duration** 12 months

# **Other Criteria**

For authorization, please submit to EviCore at evicore.com or call at 877-825-7722.

Drugs EVOTAZ

**Covered Uses** All FDA-approved indications not otherwise excluded from Health Plan

**Exclusion Criteria** 

**Required Medical Information** 

Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Plan year

Drugs EVRYSDI

# **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D.

### **Exclusion Criteria**

### **Required Medical Information**

a. Diagnosis of spinal muscular atrophy (SMA) AND b. Submission of medical records (e.g., chart notes, laboratory values) confirming the mutation or deletion of genes in chromosome 5q resulting in 1)homozygous gene deletion or mutation of SMN1 gene (e.g., homozygous deletion of exon 7 at locus 5q13)OR 2) heterozygous mutation of SMN1 gene (e.g., deletion of SMN1 exon 7 [allele 1] and mutation of SMN1 [allele 2])AND c. Patient is not dependent on either invasive ventilation or tracheostomy OR Use of non-invasive ventilation beyond use for naps and nighttime sleep AND d. Physician attests that Evrysdi is not to be initiated in a patient less than 2 month and e. Patient is not receiving concomitant chronic survival motor neuron (SMN)modifying therapy [e.g., Spinraza (nusinersen)] AND f. Patient has not previously received gene replacement therapy for the treatment of SMA [e.g., Zolgensma (onasemnogene abeparvovec-xioi)]-AND g. Submission of medical records (e.g., chart notes, laboratory values)documenting the baseline assessment

# **Age Restriction**

#### **Prescriber Restriction**

Prescribed by a neurologist with expertise in the treatment of SMA

**Coverage Duration** 3 months

#### **Other Criteria**

Reauthorization: (1) Submission of medical records (e.g., chart notes, laboratory values) with the most recent results documenting a positive clinical response to Evrysdi compared to pretreatment baseline status

#### Drugs EXJADE

# **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

Creatinine clearance less than 40 mL/minute. Platelet count less than 50 x 109/L. Poor performance status. Severe (Child-Pugh class C) hepatic impairment. High-risk myelodysplastic syndromes. Advanced malignancies. Gastrointestinal ulceration or hemorrhage.

# **Required Medical Information**

Medical documentation of FDA approved diagnosis, serum ferritin levels and serum creatinine.

# **Age Restriction**

2 years of age or older for chronic iron overload due to transfusions. 10 years of age or older for chronic iron overload due to NTDT.

**Prescriber Restriction** 

**Coverage Duration** 3 months

#### Drugs FANAPT ORAL TABLET, FANAPT ORAL TABLETS, DOSE PACK

#### **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

**Exclusion Criteria** 

**Required Medical Information** Documentation of diagnosis and treatment history.

Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Plan year

# **Other Criteria**

Approve if member has tried two following drugs: aripiprazole, clozapine, olanzapine, risperidone, quetiapine, ziprasidone

Drugs FARYDAK

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Clinically diagnosed with multiple myeloma. Trial and failure of at least two prior treatment regimens, including bortezomib and an immunomodulatory agent. Must be used in combination with bortezomib and dexamethasone.

# Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Through end of benefit year

Drugs FASENRA PEN

**Covered Uses** 

**Exclusion Criteria** 

# **Required Medical Information**

Documentation that the patient has asthma with an eosinophilic phenotype defined as blood eosinophils greater than or equal to 300 cells/µL within previous 12 months or greater than or equal to 150 cells/µL within 6 weeks of dosing AND medication will be used in combination with a corticosteroid inhaler and long acting beta2-agonist AND Patient must have experienced two or more exacerbations in the previous year OR require daily oral corticosteroids

**Age Restriction** 

**Prescriber Restriction** 

**Coverage Duration** 6 months

# Drugs

fentanyl citrate buccal lozenge on a handle

# **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

Coverage not provided in the management of acute or postoperative pain (including headache/migraines), opiod non-tolerant patients, patients with known intolerance or hypersensitivity to the drug or fentanyl.

# **Required Medical Information**

Diagnosis of cancer AND 1. Use is for breakthrough cancer pain, AND 2. Patient is opioid tolerant and taking at least 60 mg morphine/day, at least 25 mcg transdermal fentanyl/hour, at least 30 mg of oxycodone daily, at least 8 mg oral hydromorphone daily or an equianalgesic dose of another opioid for a week or longer, AND 3. Other formulary short-acting strong narcotic analgesic alternatives (other than fentanyl) have been ineffective, not tolerated, or contraindicated, AND 4. Prescriber is registered in the Transmucosal Immediate Release Fentanyl (TIRF) Risk Evaluation and Mitigation Strategy Access program.

# Age Restriction

16 years or older

# **Prescriber Restriction**

Prescribed by an oncologist or pain specialist.

**Coverage Duration** 3 months

#### **Other Criteria**

Patient must have tried and failed or not responded to the following formulary short-acting narcotics, Oxycodone and morphine. Available only to those enrolled in the Transmucosal Immediate Release Fentanyl (TIRF) EMS Program.

#### Drugs

deferiprone oral tablet 500 mg, FERRIPROX ORAL TABLET 500 MG

#### **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

Any pertinent clinical situation as defined by the product label that could affect patient safety and/or therapeutic efficacy (i.e. contraindications, warnings, precautions, adverse effects, renal or hepatic function, drug interactions, lab values, required prior or concomitant therapy, inappropriate dosing and/or duration, etc).

#### **Required Medical Information**

Diagnosis of transfusional iron overload due to thalassemia syndromes AND patient has failed prior chelation therapy with Desferal or Exjade (failure is defined as a serum ferritin level greater than 2,500 mcg/L) or patient has a contraindication or intolerance to Desferal or Exjade AND Patient has an absolute neutrophil count greater than 1.5 x 109/L.

#### Age Restriction

**Prescriber Restriction** Prescribed by a hematologist/oncologist or hepatologist

Coverage Duration Per treatment

#### Other Criteria

For renewal, patient has experienced at least a 20% reduction in serum ferritin levels and has an absolute neutrophil count greater than 0.5 x 109/L

Drugs FINTEPLA

**Covered Uses** 

**Exclusion Criteria** 

### **Required Medical Information**

Patients had a clinical diagnosis of Dravet syndrome and seizures that were inadequately controlled on at least 1 antiepileptic drug (AED) OR treatment including vagal nerve stimulation OR ketogenic diet.

Age Restriction Children 2 and older

**Prescriber Restriction** 

**Coverage Duration** Through end of benefit year

#### Drugs ICATIBANT

**Covered Uses** All FDA-approved indications not otherwise exluded for part D

# **Exclusion Criteria**

# **Required Medical Information**

Documentation of clinical diagnosis of hereditary angioedema or C1 inhibitor deficiency and having angioedema attacks.

# Age Restriction

# **Prescriber Restriction**

Must be prescribed by an allergist, immunologist, hematologist, or a physician that specializes in the treatment of HAE or related disorders.

# **Coverage Duration** 3 months

Drugs FIRDAPSE

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

**Exclusion Criteria** Member has a history of seizures

# **Required Medical Information**

The diagnosis has been confirmed by one of the following: A) Presence of anti-P/Q-type voltage-gated calcium channel (VGCC) antibodies OR B) Characteristic electromyography (EMG).

Age Restriction

Prescriber Restriction Neurologist

**Coverage Duration** Through the benefit year

# Fondaparinux (ARIXTRA)

#### Drugs

fondaparinux subcutaneous syringe 10 mg/0.8 mL, 2.5 mg/0.5 mL, 5 mg/0.4 mL, 7.5 mg/0.6 mL

#### Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

CrCl (EGFR) less than 30mL/min. Patient's weight less than 50kg.

#### **Required Medical Information**

Clinical documentation of FDA approved indication for treatment. Patient's weight and creatinine clearance (CrCl).

# Age Restriction

18 years and older.

# **Prescriber Restriction**

# **Coverage Duration**

Per treatment. Post-op DVT prophylaxis 1. hip/knee replacement max of 35 days. 2. abdominal surgery

#### Drugs FORTEO SUBCUTANEOUS PEN INJECTOR 20 MCG/DOSE (600MCG/2.4ML)

# **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

Because of an increased incidence of osteosarcoma, Forteo should not be prescribed for patients who are at increased baseline risk for osteosarcoma (including those with Paget's disease of bone or unexplained elevations of alkaline phosphatase, open epiphyses, or prior radiation therapy involving the skeleton).

# **Required Medical Information**

Approve the following: 1. Treatment of postmenopausal women with osteoporosis at high risk for fracture OR 2. Increase of bone mass in men with primary or hypogonadal osteoporosis at high risk for fracture, AND member meets the following criteria: a. fracture OR b. BMD screening results of -2.5 or below OR c. previously failure / contraindication / intolerance of an oral bisphosphonate AND Prolia 3. Treatment of men and women with osteoporosis associated with sustained systemic glucocorticoid therapy at high risk for fracture, AND member meets the following criteria: a. previous fracture, OR b. multiple risk factors for fracture, OR c. previous failure/contraindication/intolerance of the following: oral bisphosphonate AND Prolia. Patient has not received more than 2 years of therapy with Forteo.

# Age Restriction

# **Prescriber Restriction**

# **Coverage Duration**

For initial therapy up to 1 year. Continuation up to 1 year not to exceed 2 years of total therapy.

# **Other Criteria**

Treatment failure is defined as documented continued bone loss after at leaast three months despite treatment with a bisphosphonate or Prolia. Note: Since the effects of long-term treatment with teriparatide are not known at this time, therapy for more than 2 years duration is considered experimental and investigational.

#### Drugs FOTIVDA

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Diagnosis of Renal cell carcinoma, Advanced, relapsed or refractory following 2 or more prior systemic therapies

Age Restriction 18 years or older

Prescriber Restriction Oncologist

# **Coverage Duration**

End of Benefit Year

# **Other Criteria**

1) Must have tried and failed or have a contraindication to other NCCN recommended therapies 2) Must have an Eastern Cooperative Oncology Group (ECOG) performance status of 0 or 1

# Drugs FUZEON SUBCUTANEOUS RECON SOLN

### **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

# **Required Medical Information** Clinical documentation of FDA approved indication for treatment.

# Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Plan year

**Other Criteria** For treatment-experienced patients.

#### Drugs FYCOMPA ORAL SUSPENSION, FYCOMPA ORAL TABLET 10 MG, 12 MG, 2 MG, 4 MG, 6 MG, 8 MG

# **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D. All FDA-approved indications not otherwise excluded from Health Plan.

**Exclusion Criteria** 

**Required Medical Information** 

Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Plan year

# Drugs GAVRETO

**Covered Uses** 

**Exclusion Criteria** 

# **Required Medical Information**

Patient has a diagnosis of metastatic non-small cell lung cancer (NSCLC) AND Patient's disease is RET fusion-positive detected by an FDA approved test.

Age Restriction

**Prescriber Restriction** 

**Coverage Duration** 3 months

Drugs GENVOYA

**Covered Uses** All FDA-approved indications not otherwise excluded from Health Plan

**Exclusion Criteria** 

**Required Medical Information** 

**Age Restriction** 

**Prescriber Restriction** 

**Coverage Duration** Plan year

Drugs GILENYA

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

# **Required Medical Information**

Diagnosis of relapsing forms of multiple sclerosis and must have tried and failed one generic alternative agent such as dimethyl fumarate or glatiramer acetate.

# Age Restriction

**Prescriber Restriction** Prescribed by or in consultation with a neurologist or MS specialist.

#### **Coverage Duration** Plan year

Drugs GILOTRIF

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Documented diagnosis of 1) Metastatic non-small cell lung cancer (NSCLC) with tumors that have non-resistant epidermal growth factor receptor (EGFR) mutations as detected by an FDA-approved test OR 2) Metastatic squamous non-small cell lung cancer progressing after platinum-based chemotherapy.

### **Age Restriction**

**Prescriber Restriction** 

**Coverage Duration** Through end of benefit year

#### GLEOSTINE

#### Drugs GLEOSTINE

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

**Exclusion Criteria** 

**Required Medical Information** Must be prescribed for an FDA approved indication.

### Age Restriction

Prescriber Restriction Oncologist

**Coverage Duration** 12 months

#### Drugs OMNITROPE

#### **Covered Uses**

All FDA approved indications not otherwise excluded by Health Plan. Additional off-label coverage is provided for (note - some growth hormone drugs may be labeled for 1 or more of these indications): adult growth hormone deficiency, growth failure in children small for gestational age or with intrauterine growth retardation, idiopathic short stature, GH deficiency associated with Turner Syndrome, growth failure secondary to chronic renal failure/insufficiency in children who have not received a renal transplant, short stature associated with Noonan Syndrome, short bowel syndrome, and for the treatment of Prader-Willi Syndrome.

#### **Exclusion Criteria**

Coverage is not provided for constitutional delayed growth

#### **Required Medical Information**

Pediatric GHD: epiphyses must be confirmed open in patients 10 years of age and older, AND 1. diagnosis confirmed by any 2 provocative tests or by both low IGF-1 and IGFBP-3 levels in patients who meet the height related conditions of coverage, 2. diagnosis confirmed by 2 provocative tests and both low IGF-1 and IGF-BP3 in patients not meeting height related coverage conditions, or 3. 3 pituitary hormone deficiencies in pt with irreversible hypothalamic-pituitary structural lesions or panhypopituitarism. Growth failure from CRF: PGHD criteria must be met without the provocative tests or IGF-1 and IGF-BP3 related conditions. Idiopathic Short Stature: epiphyses must be confirmed as open in patients greater than or equal 10 years of age, height must be less than or equal - 2.25 sds from the mean. Small for Gestational Age: failure to manifest catch up growth by age 2 defined as birth weight, birth length, or both that are more than 2 sds mean normal values following adjustment for age and gender. Turner's syndrome and Noonan Syndrome: epiphyses must be confirmed as open and when on therapy. Adult GHD: requires either 1. a negative GH provocative test when the AGHD is due to childhood onset GHD, pituitary or hypothalamic disease, surgery or radiation therapy, or trauma, OR 2. 3 pituitary hormone deficiencies and baseline serum IGF-I levels below the age- and sex-appropriate reference range when the AGHD is due to irreversible hypothalamic-pituitary structural lesions are from childhood. Short bowel syndrome: when receiving specialized nutritional support.

#### Age Restriction

7 years of age or older for Idiopathic short stature

#### **Prescriber Restriction**

Pediatric endocrinologist for ISS

#### **Coverage Duration**

1 month for short bowel syndrome, 12 months for other indications

#### **Other Criteria**

Height related conditions of coverage: 1. height below the third percentile for their age and gender related height, 2. growth velocity subnormal greater than or equal 2 standard deviations (sds) from the age related mean, 3. delayed skeletal maturation greater than or equal 2 sds below the age/gender related mean. Renewals for PGHD, CFR, SGA, Turner's and Noonan Syndromes require growth response of greater than or equal 4.5 cm/yr (pre-pubertal) or greater than or equal 2.5 cm/yr (post-pubertal) AND open epiphyses. For pediatric patients with irreversible hypothalamic-pituitary structural lesions or panhypopituitarism coverage is renewable if the patient has had 3 pituitary hormone deficiencies. Renewals for short bowel syndrome is provided in the presence of clinical benefit (such as, decreasing the patient?s intravenous nutritional requirements). Renewals for Prader-Willi syndrome is provided if pt has increase in lean body mass or decrease in fat mass. Renewals for AGHD is provided in the presence of a growth response of greater than or equal 1.5 cm/yr AND open epiphyses. Renewals for AGHD is provided in the presence of clinical benefit (e.g., increase in total lean body mass, increase in IGF-1 and IGFBP-3 levels, or increase in exercise capacity).

#### Drugs HARVONI ORAL TABLET 90-400 MG

# **Covered Uses**

All FDA-approved indications not otherwise excluded from Health Plan.

# **Exclusion Criteria**

1. Active IV drug users, 2. Active alcohol users, 3. Reinfection after 6 months cure

### **Required Medical Information**

For initial authorization (12 weeks maximum), provider must submit completed HCV Tx form, medical records documenting the diagnosis of chronic hepatitis C with genotype and subtype, if applicable, medical records documenting viral load taken within 6 months of beginning therapy AND submit medical records documenting advanced fibrosis as corresponding to a FibroSure or a Liver Biopsy proven. Other fibrosis scores, physical findings, or clinical evidence consistent with cirrhosis as attested by the prescribing physician may be also considered. For any retreatment or extension of PA, 100% compliance will be required (Claim hx). Urine alcohol metabolite and drug screen required.

### Age Restriction

Patient must be 18 years of age or older

#### **Prescriber Restriction**

Prescribed by, a gastroenterologist, hepatologist, or infectious disease physician.

### **Coverage Duration**

Based on the AASLD treatment guidelines

#### **Other Criteria**

Criteria and coverage durations will be applied consistent with current AASLD/IDSA guidance. Documentation of F2-F4 fibrosis (fibrosis score of 0.48 and greater) or patient is Health Care worker in direct patient care setting. Must have contraindication to or be unable to tolerate Mavyret, generic Ledipasvir/Sofosbuvir, and generic Sofosbuvir/Velpatasvir.

#### Drugs HERCEPTIN HYLECTA

#### Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

#### **Exclusion Criteria**

#### **Required Medical Information**

Documentation of 1) HER2 overexpressing node positive or node negative breast cancer as part of a treatment regimen consisting of a) doxorubicin, cyclophosphamide, and either paclitaxel or docetaxel or b)docetaxel and carboplatin or c) as a single agent following multi-modality anthracycline based therapy OR 2)HER2-overexpressing metastatic breast cancer either a) in combination with paclitaxel for first-line treatment or b) as a single agent for patients who have received one or more chemotherapy regimens for metastatic disease.

#### Age Restriction

Prescriber Restriction Oncology

**Coverage Duration** 3 months

Drugs HORIZANT

**Covered Uses** All FDA-approved indications not otherwise excluded from Health Plan

#### **Exclusion Criteria**

#### **Required Medical Information**

Documented diagnosis of postherpetic neuralgia (PHN) and trial of gabapentin, or diagnosis of moderate to severe primary restless legs syndrome (RLS) in adults and trial of ropinirole and pramipexole.

Age Restriction

**Prescriber Restriction** 

Coverage Duration Plan year

#### Drugs

#### HUMIRA PEN, HUMIRA PEN CROHNS-UC-HS START, HUMIRA PEN PSOR-UVEITS-ADOL HS, HUMIRA SUBCUTANEOUS SYRINGE KIT 40 MG/0.8 ML, HUMIRA(CF), HUMIRA(CF) PEDI CROHNS STARTER SUBCUTANEOUS SYRINGE KIT 80 MG/0.8 ML, 80 MG/0.8 ML-40 MG/0.4 ML, HUMIRA(CF) PEN, HUMIRA(CF) PEN CROHNS-UC-HS, HUMIRA(CF) PEN PSOR-UV-ADOL HS

#### **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D. All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

Using in combination with other TNF antagonists, IL-1R antagonists, janus kinase inhibitor, anti-cd20 monoclonalantibodies or selective co-stimulation modulators.

#### Required Medical Information

Diagnosis of 1 of the following: A) Mod-severe rheumatoid arthritis and trial of 1 or more non-biologic disease modifying anti-rheumatic drugs (DMARD) (e.g., hydroxychloroquine [HCQ], sulfasalazine, methotrexate [MTX], leflunomide, azathioprine, cyclosporine) for at least 3 months, or B) Mod-severe polyarticular juvenile idiopathic arthritis (JIA) and trial of 1 or more non-biologic DMARDs (e.g., HCQ, sulfasalazine, MTX, leflunomide, azathioprine, cyclosporine) for at least 3 months, or C) Psoriatic arthritis and trial of MTX, or D) Ankylosing spondylitis and trial of 1 or more NSAIDs, or E) Modsevere chronic plaque psoriasis (affecting more than 5% of body surface area or crucial body areas such as the hands, feet, face, or genitals) and trial of at least 1 of the following: phototherapy (including but not limited to Ultraviolet A with a psoralen [PUVA] and/or retinoids [RePUVA] for at least 1 continuous month or 1 or more oral systemic treatments (e.g., MTX, cyclosporine, acitretin, sulfasalazine) for at least 3 months, or F) Mod-severe Crohn's disease and trial of 2 or more of the following: corticosteroids (e.g., prednisone, methylprednisolone) or non-biologic DMARDs (e.g., azathioprine. MTX. mercaptopurine), or G) Mod-severe ulcerative colitis trial of 2 or more of the following: corticosteroids (e.g., prednisone, methylprednisolone), 5-ASA (i.e. mesalamine, sulfasalazine, balsalazide, olsalazine) or non-biologic DMARDs (azathioprine, MTX, mercaptopurine), or H) Hidradenitis suppurativa, or I) Non-infectious intermediate, posterior, and panuveitis and trial of 1 or more of the following: periocular, intraocular, or systemic corticosteroids, immunosuppressants (azathioprine, MTX, mycophenolate mofetil, cyclophosphamide, cyclosporine). A trial is defined as an inadequate response, intolerance or contraindication to the therapy.

Age Restriction

**Prescriber Restriction** 

**Coverage Duration** 12 months

Drugs HYCAMTIN

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

**Exclusion Criteria** 

**Required Medical Information** Must be prescribed for an FDA approved indication

Age Restriction

Prescriber Restriction Oncologist

**Coverage Duration** 12 months

Drugs IBRANCE

**Covered Uses** All FDA-approved indications not otherwise excluded from Health Plan.

**Exclusion Criteria** 

**Required Medical Information** 

Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Plan year

#### Drugs IDAMYCIN PFS

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

**Exclusion Criteria** 

**Required Medical Information** Must be prescribed for an FDA approved indication

# Age Restriction

Prescriber Restriction Oncologist

**Coverage Duration** 12 months

Drugs IDHIFA

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

### **Required Medical Information**

1. Diagnosis of relapsed or refractory acute myeloid leukemia (AML) with an isocitrate dehydrogenase-2 (IDH2) mutation AND 2. Prescriber agrees to monitor for signs and symptoms of differentiation syndrome.

### Age Restriction

**Prescriber Restriction** Oncologist or Hematologist

**Coverage Duration** Through the end of benefit year

#### Drugs

imatinib oral tablet 100 mg, 400 mg

#### **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

#### **Required Medical Information**

Diagnosis of one of the following: A) Philadelphia chromosome-positive chronic myelogenous leukemia (Ph+ CML), or B) Ph+ acute lymphoblastic leukemia (ALL), or C) Gastrointestinal tumor (GIST) where patient has documented c-KIT (CD117) positive unresectable or metastatic malignant GIST or patient had resection of c-KIT positive GIST and imatinib will be used as an adjuvant therapy, or D) Dermatofibrosarcoma protuberans that is unresectable, recurrent, or metastatic, or E) Hypereosinophilic syndrome or chronic eosinophilic leukemia, or F) Myelodysplastic syndrome or myeloproliferative disease associated with platelet-derived growth factor receptor gene re-arrangements, or G) Aggressive systemic mastocytosis without the D816V c-KIT mutation or with c-KIT mutation or with c-KIT mutational status unknown.

#### Age Restriction

1 year of age or older - newly diagnosed CML in the chronic phase or newly diagnosed Ph+ ALL. 18 years of age or older for other indications.

**Prescriber Restriction** 

Coverage Duration Plan year

# Drugs IMBRUVICA ORAL CAPSULE 140 MG, 70 MG, IMBRUVICA ORAL TABLET

Covered Uses

All FDA-approved indications not otherwise excluded from Health Plan.

**Exclusion Criteria** 

**Required Medical Information** 

**Age Restriction** 

**Prescriber Restriction** 

**Coverage Duration** 3 Months

Drugs INBRIJA

# Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

#### **Exclusion Criteria**

patient is not currently taking or has recently (within 2 weeks) taken a nonselective monoamine oxidase (MAO) inhibitor (e.g., phenelzine and tranylcypromine)

#### **Required Medical Information**

Patient is currently treated with carbidopa/levodopa AND is experiencing intermittent OFF episodes secondary to Parkinson's disease

Age Restriction

**Prescriber Restriction** 

**Coverage Duration** 3 months

**Other Criteria** 

Reauthorization requires physician attestation of medications efficacy

Drugs ICLUSIG ORAL TABLET 15 MG, 45 MG

**Covered Uses** 

**Exclusion Criteria** 

#### **Required Medical Information**

Documented diagnosis of one of the following: 1. T315I-positive chronic myeloid leukemia (CML)OR 2. T315I-positive Philadelphia chromosome positive acute lymphoblastic leukemia (Ph+ ALL)OR 3. Chronic phase (CP) chronic myeloid leukemia (CML)with resistance or intolerance to at least two prior tyrosine kinase inhibitors OR 4. Accelerated phase (AP) or blast phase (BP) chronic myeloid leukemia (CML)AND no other tyrosine kinase (TKI) therapy is indicated OR 5. Philadelphia chromosome positive acute lymphoblastic leukemia (Ph+ ALL)AND no other tyrosine kinase (TKI) therapy is indicated or 5.

Age Restriction 18 years of age or older

**Prescriber Restriction** 

**Coverage Duration** 3 months

#### Drugs INCRELEX

#### **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

Insulin-like growth factor therapy is considered NOT medically necessary when any of the following criteria are met: Final adult height has been reached as determined by the 5th percentile of adult height OR the bone epiphyses are closed OR the patient is older than 18 years of age. Contraindicated in neonates, patients with closed epiphyses, and suspected neoplasia.

#### **Required Medical Information**

1. All of the following: a. Diagnosis of severe primary IGF-1 deficiency. b. Height standard deviation score of -3.0 or less. c. Basal IGF-1 standard deviation score of -3.0 or less. d. Normal or elevated growth hormone. e. Open finger epiphyses on last boneradiograph GH gene deletion: a. Diagnosis of growth hormone gene deletion who have developed neutralizing antibodies to GH, AND b. Have open finger epiphyses on last bone radiograph.

### Age Restriction

The patient is between 2 years -18 years old for Increlex therapy

# Prescriber Restriction

Must be endocrinologist to prescribe

**Coverage Duration** 6 months to 1 year

#### **Other Criteria**

Not a substitute for GH treatment. For renewal, Patient had a minimum growth rate of at least 2 cm/year.

Drugs INLYTA

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

#### **Exclusion Criteria**

#### **Required Medical Information**

Diagnosis of 1) advanced renal cell carcinoma AND patient failed one or more systemic therapies for renal cell carcinoma (e.g., sunitinib-, bevacizumab-, temsirolimus-, or cytokine-containing regimens) OR 2) advanced renal cell carcinoma and the medication will be used in combination with avelumab OR 3) advanced renal cell carcinoma (RCC) and the medication will be used in combination with pembrolizumab.

Age Restriction 18 years of age or older

**Prescriber Restriction** 

**Coverage Duration** 3 months

Drugs INQOVI

**Covered Uses** 

**Exclusion Criteria** 

#### **Required Medical Information**

Patient has a diagnosis of myelodysplastic syndromes (MDS), including previously treated and untreated, de novo and secondary MDS with the following French-American-British subtypes (refractory anemia, refractory anemia with ringed sideroblasts, refractory anemia with excess blasts, and chronic myelomonocytic leukemia [CMML]) and intermediate-1, intermediate-2, and high-risk International Prognostic Scoring System groups AND patient is not receiving Inqovi concomitantly with intravenous decitabine AND patient has tried and failed or has a contraindication to SQ azacitadine.

**Age Restriction** 

**Prescriber Restriction** 

**Coverage Duration** 3 months

#### INREBIC

#### Drugs INREBIC

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

#### **Exclusion Criteria**

Patients on treatment with ruxolitinib before initiation must taper and discontinue according to ruxolitinib prescribing information

#### **Required Medical Information**

Must provide labs showing patient is not thiamine deficient before starting drug

Age Restriction 18 years of age and older

**Prescriber Restriction** Oncologist or Hematologist

**Coverage Duration** 3 months

#### Drugs INTELENCE ORAL TABLET 100 MG, 200 MG, 25 MG

#### **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

#### **Required Medical Information** Clinical documentation of FDA approved indication for treatment.

#### **Age Restriction**

**Prescriber Restriction** 

**Coverage Duration** Plan year

**Other Criteria** For treatment-experienced patients.

#### Drugs INTRON A INJECTION RECON SOLN

# **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

#### **Required Medical Information**

Type B viral Hepatitis (HBeAg positive): Serum HBsAg positive for at least six months, AND elevated serum ALT 2 times ULN or moderate to severe hepatitis or fibrosis on biopsy. Type B Viral Hepatitis (HBeAg negative) HBsAG positive for at least 6 months AND BHV DNA level of 2000 IU/ml or more than 11,200 copies/ml AND One of the following, persistent ALT 2 times UNL or moderate to severe hepatitis or fibrosis on biopsy. Documentation must be provided showing trial and failure to our preferred agent Peg-Intron. Chronic Hepatitis C: Positive HCV antibody and HCV RNA. Documentation must be provided showing trial and failure to our preferred agent Peg-Intron. Condyloma Acuminatum or Perianal Warts: Must have documentation of trial and failure to preferred alternative or intolerance/contraindication to preferred alternatives. For external perianal warts, condylox gel, for external genital warts, podofilox, or imiquimod. Hairy Cell Leukemia: Medical documentation indicating diagnosis. Malignant Melanoma: Indicated as adjuvant to surgical treatment with malignant melanoma who are free of disease but at high risk for systemic recurrence, within 56 days of surgery. Follicular Lymphoma: Indicated for the initial treatment of clinically aggressive follicular Non-Hodgkins Lymphoma in conjunction with anthracycline-containing combination chemotherapy. Efficacy in patients with low-grade, low-tumor burden follicular Non-Hodgkins Lymphoma has not been demonstrated. AIDS-Related Kaposis Sarcoma: Indicated for the treatment of selected patients. The likelihood of response to therapy is greater in patients who are without systemic symptoms, who have limited lymphadenopathy and who have a relatively intact immune system as indicated by total CD4 count.

#### **Age Restriction**

For Hepatitis B- age 1 or older, For Hepatitis C - age 3 or older, All other diagnoses- 18 years or older.

**Prescriber Restriction** 

Coverage Duration Plan year

Drugs INVEGA HAFYERA, INVEGA SUSTENNA, paliperidone oral tablet extended release 24hr 1.5 mg, 3 mg, 6 mg, 9 mg

#### **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

#### **Required Medical Information**

Diagnosis of Schizophrenia AND documented treatment failure or intolerable side effects from treatment with two formulary antipsychotic medications such as risperidone, ziprasidone,

#### Age Restriction

**Prescriber Restriction** 

Coverage Duration Plan year

#### Drugs INVIRASE ORAL TABLET

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

**Required Medical Information** Clinical documentation of FDA approved indication for treatment.

### Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Plan year

# **Other Criteria**

Pretreatment EKG should be done and therapy is NOT to be initiated if QT interval exceeds 450 msec. Must be used in combination with ritonavir.

#### Drugs IRESSA

**Covered Uses** All FDA-approved indication not otherwise excluded from Part D.

# **Exclusion Criteria**

**Required Medical Information** Diagnosis

# Age Restriction

**Prescriber Restriction** Must be prescribed by Oncologist

**Coverage Duration** 3 mos initial, renewable in 6 month increments

#### Drugs ISENTRESS HD, ISENTRESS ORAL POWDER IN PACKET, ISENTRESS ORAL TABLET, ISENTRESS ORAL TABLET, CHEWABLE

#### **Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

**Required Medical Information** Clinical documentation of FDA approved indication for treatment.

#### **Age Restriction**

**Prescriber Restriction** 

**Coverage Duration** Plan year

#### Itraconazole (Sporanox)

Drugs

itraconazole oral capsule

**Covered Uses** 

All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

**Required Medical Information** 

#### Age Restriction

**Prescriber Restriction** 

# **Coverage Duration** 3 months

**Other Criteria** 

For onychomycosis - must have documented failure, intolerance or contraindication to terbinafine.

#### Drugs GAMMAGARD LIQUID

#### **Covered Uses**

All medically accepted indications not otherwise excluded by Health Plan, chronic inflammatory demyelinating polyneuropathy, multifocal motor neuropathy, dermatomyositis, polymyositis, Guillain-Barre syndrome (GBS), relapsing-remitting multiple sclerosis (RRMS), myasthenia gravis, Lambert-Eaton myasthenic syndrome, Kawasaki syndrome, idiopathic thrombocytopenic purpura, pure red cell aplasia (PRCA), fetal/neonatal alloimmune thrombocytopenia, and prophylaxis of bacterial infections in B-cell chronic lymphocytic leukemia (CLL), bone marrow/hematopoietic stem cell transplant (BMT/HSCT) recipients, and pediatric HIV infection.

#### **Exclusion Criteria**

IgA deficiency with antibodies to IgA and a history of hypersensitivity. History of anaphylaxis or severe systemic reaction to human immune globulin or product components.

#### **Required Medical Information**

Documentation of diagnosis and previous treatment. For dermatomyositis and polymyositis: standard 1st line treatments (corticosteroids or immunosuppressants) have been tried but were unsuccessful or not tolerated OR patient is unable to receive standard therapy because of a contraindication or other clinical reason. For GBS: physical mobility must be severely affected such that the patient requires an aid to walk AND IVIG therapy must be initiated within 2 weeks of symptom onset. For RRMS: standard 1st line treatments (e.g. interferon, glatiramer, dimethyl fumarate) have been tried but were unsuccessful or not tolerated OR patient is unable to receive standard therapy because of a contraindication or other clinical reason. For CLL: serum IgG less than 500 mg/dL OR a history of recurrent bacterial infections. For BMT/HSCT: serum IgG less than 400 mg/dL. For pediatric HIV infection: serum IgG less than 400 mg/dL OR a history of recurrent bacterial infections. PRCA is secondary to parvovirus B19 infection. For all indications: patients with any of the following risk factors for renal dysfunction must receive the minimum dose or concentration available of IVIG and the minimum infusion rate practicable: pre-existing renal insufficiency, diabetes mellitus, age over 65 years, volume depletion, sepsis, paraproteinemia, or receiving concomitant nephrotoxic drugs. For all indications: patients with any of the following risk factors for thrombosis must receive the minimum dose or concentration available of IVIG and the minimum infusion rate practicable: pre-existing renal insufficiency, diabetes mellitus, age over 65 years, volume depletion, sepsis, paraproteinemia, or receiving concomitant nephrotoxic drugs. For all indications: patients with any of the following risk factors for thrombosis must receive the minimum dose or concentration available of IVIG and the minimum infusion rate practicable: age 45 years or older, prolonged immobilization, hypercoagulable conditions, history of venous or arterial thrombosis, use of

Age Restriction

**Prescriber Restriction** 

Coverage Duration Plan year

#### Drugs JAKAFI ORAL TABLET 10 MG, 15 MG, 20 MG, 25 MG, 5 MG

# **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

#### **Required Medical Information**

Statement of diagnosis for treatment of patients with intermediate or highrisk myelofibrosis, including primary myelofibrosis, post-polycythemia vera myelofibrosis and post-essential thrombocythemia myelofibrosis and lab work indicating a complete blood count and platelet count before initiating therapy and recent lab work indicating complete blood count and platelet count for a dosage adjustment. Lab work must indicate platelets are more than 50 x 109/L and dose must be less than 50 mg per day. No dose increases will be approved within 4 weeks of therapy and not more frequently than every 2 weeks. If no spleen reduction or symptom improvement after 6 months then discontinue the drug.

#### Age Restriction

18 years or older

#### **Prescriber Restriction**

Myelofibrosis: Prescribed by a Hematologist/Oncologist

**Coverage Duration** 3 months

#### Drugs JYNARQUE ORAL TABLETS, SEQUENTIAL 45 MG (AM)/ 15 MG (PM), 60 MG (AM)/ 30 MG (PM), 90 MG (AM)/ 30 MG (PM)

#### **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D.

#### **Exclusion Criteria**

#### **Required Medical Information**

Medication requested is being used to slow kidney function decline AND Liver function laboratory values (ALT, AST and bilirubin) have been reviewed and are appropriate before initiation.

#### Age Restriction

Patient is 18 years of age or older

#### **Prescriber Restriction**

#### Coverage Duration

Initial Authorization will be for 3 months. Reauthorization will be for 1 year.

Drugs KALETRA

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

**Exclusion Criteria** 

**Required Medical Information** Clinical documentation of FDA approved indication for treatment.

Age Restriction

**Prescriber Restriction** 

Coverage Duration Plan year

#### Drugs KALYDECO ORAL GRANULES IN PACKET 50 MG, 75 MG, KALYDECO ORAL TABLET

#### **Covered Uses**

All FDA-approved indications not otherwise excluded from Health Plan.

#### **Exclusion Criteria**

#### **Required Medical Information**

Medical documentation of cystic fibrosis AND member has a G551D, G1244E, G1349D, G178R, G551S, R117H, S1251N, S1255P, S549N, or S549R mutation in the CFTR gene AND member does not have a Homozygous F508del mutation in CFTR gene.

#### Age Restriction

Ivacaftor oral granules are approved in patients 2 years of age and older. Ivacaftor oral tablets are approved in patients 6 years of age and older.

# Prescriber Restriction

Endocrinologist or Pulmonologist

Coverage Duration Plan year

### Drugs KESIMPTA PEN

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

### **Required Medical Information**

Diagnosis of relapsing forms of multiple sclerosis and must have tried and failed one generic alternative agent such as dimethyl fumarate or glatiramer acetate

### Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Through end of benefit year

# Drugs KISQALI, KISQALI FEMARA CO-PACK

#### **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D.

### **Exclusion Criteria**

#### **Required Medical Information** Documentation that the member has tried and failed or has a contraindication to Ibrance.

# Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Through end of benefit year

# Drugs

#### KUVAN ORAL TABLET, SOLUBLE, sapropterin oral tablet, soluble

#### **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

#### **Required Medical Information**

#### Age Restriction 1 month and older

#### **Prescriber Restriction**

#### **Coverage Duration**

Initial: 2 months. Renewal: through plan year

#### **Other Criteria**

For initial approval, Patient will have phenylalanine levels measured one week after starting therapy and periodically for up to two months of therapy to determine response. For renewal, patient has been determined to be a responder to therapy (i.e. phenylalanine levels have decreased by at least 30% from baseline) and phenylalanine levels will be measured periodically during therapy.

#### Lamivudine (EPIVIR)

#### Drugs

lamivudine oral solution, lamivudine oral tablet 100 mg, 150 mg, 300 mg

# **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

**Required Medical Information** Clinical documentation of FDA approved indication for treatment.

#### **Age Restriction**

**Prescriber Restriction** 

Coverage Duration Plan year

### LATUDA

### Drugs LATUDA ORAL TABLET 120 MG, 20 MG, 40 MG, 60 MG, 80 MG

#### Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

### **Exclusion Criteria**

#### **Required Medical Information** Trial of at least two other drugs: lurasidone, risperidone, questiapine, or cariprazine.

#### **Age Restriction**

**Prescriber Restriction** 

#### **Coverage Duration** Through the end of benefit year

#### Drugs LEDIPASVIR-SOFOSBUVIR

#### **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D.

#### **Exclusion Criteria**

#### **Required Medical Information**

Provider must submit medical records documenting the following: 1) medical diagnosis of Chronic Hepatitis C with labs documenting genotype and subtype, AND 2) medical records documenting viral load taken within 6 months of beginning therapy, AND 3) fibrosis score to confirm appropriate duration of treatment, AND 4) documentation of previous HCV therapies to confirm appropriate duration of treatment.

### Age Restriction

Patient must be 12 years of age or older

#### **Prescriber Restriction**

Prescribed by, a gastroenterologist, hepatologist, or infectious disease physician.

#### **Coverage Duration**

12 to 24 weeks based on the AASLD treatment guidelines

#### **Other Criteria**

Criteria and coverage durations will be applied consistent with current AASLD/IDSA guidance.

Drugs LENVIMA

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

**Exclusion Criteria** 

**Required Medical Information** 

Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Through benefit year

#### Drugs

ambrisentan, LETAIRIS

#### **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

Known or suspected pregnancy. Treat women of child-bearing potential only after a negative pregnancy test and treat only women who are using two reliable methods of contraception OR have had a tubal sterilization OR a Copper T 380A IUD or LNg 20 IUD inserted.

#### **Required Medical Information**

Diagnosis of Pulmonary Arterial Hypertension (PAH) AND pregnancy must be excluded prior to the start of therapy and will be prevented thereafter with two forms of reliable contraception in female patients of reproductive potential. Trial and failure of Revatio or Adcirca.

Age Restriction 18 years and older

**Prescriber Restriction** 

Coverage Duration 3 months

#### Drugs LEUKINE INJECTION RECON SOLN

# **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

Chemotherapy or radiotherapy within 24 hours or concomitantly, excess leukemic myeloid blasts in the bone marrow or blood (10% or greater), hypersensitivity to granulocyte-macrophage colony-stimulating factor (GM-CSF) or yeast-derived products, allergic or anaphylactoid reactions to the medication in the past.

# **Required Medical Information**

Medical statement indicating diagnosis AND trial and failure of preferred agent neupogen AND Absolute Neutrophil Count less than 10,000/mm3 and CBC with differential.

# Age Restriction

Patients requiring prophylaxis of febirle neutropenia in acute myelogenous leukemia following induction chemotherapy must be at least 55 years of age, other diagnoses do not specify an age restriction

Prescriber Restriction Oncologist or Hematologist

Coverage Duration Plan year

#### Drugs

leuprolide

# **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

Pregnancy in patients with child-bearing potential. Breastfeeding. Undiagnosed abnormal vaginal bleeding.

# **Required Medical Information**

Diagnosis of one of the following: A) Advanced or metastatic prostate cancer (7.5 mg 1-month, 22.5 mg 3-month, 30 mg 4-month, & 45 mg 6-month depots only), or B) Central precocious puberty (Lupron Depot-Ped) AND submission of pubertal gonadal sex steroid levels (testosterone greater than 30 ng/dL, estradiol greater than 20 pg/mL AND a pubertal LH increase upon native GnRH stimulation AND pelvic ultrasound assessment (girls) is required for approval along with notes indicating premature development of secondary sexual characteristics at or before the age of 8 yrs in girls and 9 yrs in boys and significant advancement of bone age and/or a poor adult height prediction AND other causes of sexual precocity must be excluded.

Age Restriction

**Prescriber Restriction** 

Coverage Duration Plan year

#### Drugs

fosamprenavir, LEXIVA

#### **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

Any pertinent clinical situation as defined by the product label that could affect patient safety and/or therapeutic efficacy (i.e. contraindications, warnings, precautions, adverse effects, renal or hepatic function, drug interactions, lab values, required prior or concomitant therapy, inappropriate dosing and/or duration, etc).

# **Required Medical Information**

Clinical documentation of FDA approved indication for treatment.

Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Plan year

# LIDOCAINE PRODUCTS

#### Drugs

lidocaine topical adhesive patch, medicated 5 %, lidocaine topical ointment

#### **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D. All FDA-approved indications not otherwise excluded by Health Plan

**Exclusion Criteria** 

**Required Medical Information** 

Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Plan year

# Drugs

linezolid

# **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

Patients that are currently myelosuppressed due to any cause

#### **Required Medical Information**

Culture and sensitivity reports verifying: 1. VRE infections within past 30 days. 2. Nosocomial pneumonia (MRSA) within past 30 days. 3. Nosocomial or CAP (MSSA or S. pneumoniae) within past 30 days and failure/resistance to 2 preferred antibiotics 4. Complicated SSI without osteomyelitis (MRSA) within past 30 days. 5. Uncomplicated SSI (MRSA) within past 30 days or empirical treatment of uncomplicated or community-acquired complicated SSI without osteomyelitis (MRSA likely) and failure/resistance to 2 preferred antibiotics. 6. Uncomplicated or complicated SSI without osteomyelitis (MSSA, S. pyogenes, or S. agalactiae (complicated SSI only)) within past 30 days and failure/resistance to 2 preferred antibiotics.

# **Age Restriction**

# **Prescriber Restriction**

Presribing physician must be an infectious disease specialist

# **Coverage Duration**

Non-MRSA nosocomial or community acquired pneumonia, SSI: 14 days Other uses: 28 days

#### **Other Criteria**

Nosocomial or community acquired pneumonia (MSSA or S. pneumoniae) preferred antibiotics: Amoxicilin/Clavulanate, Azithromycin, Cephalexin, Clarithromycin, Levaquin. Uncomplicated SSI (MRSA) or empirical treatment of patients with uncomplicated or community-acquired complicated SSI without osteomyelitis (MRSA likely) preferred antibiotics: Trimethoprim/sulfamethoxazole, Tetracycline, Doxycycline, Minocycline, Clindamycin. Uncomplicated or complicated SSI without osteomyelitis (MSSA, S. pyogenes, or S. agalactiae (complicated SSI only)) preferred antibiotics: Amoxicillin/clavulanate, Cephalexin, Ciprofloxacin, Clindamycin, Levaguin, Trimethoprim/Sulfamethoxazole, Dicloxacillin. Drugs LOKELMA

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

#### **Required Medical Information**

Documentation of elevated serum potassium (greater than 5.0 mEq/L) and the beneficiary has failure, contraindication or intolerance to sodium polystyrene sulfonate oral suspension.

Age Restriction 18 years and older

**Prescriber Restriction** 

**Coverage Duration** 6 months

Drugs LONSURF

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Documented diagnosis of 1) metastatic colorectal cancer previously treated with fluoropyrimidine-, oxaliplatin- and irinotecan-based chemotherapy, an anti-VEGF biological therapy, and if RAS wild-type, an anti-EGFR therapy OR 2) metastatic gastric or gastroesophageal junction adenocarcinoma previously treated with at least two prior lines of chemotherapy that included a fluoropyrimidine, a platinum, either a taxane or irinotecan, and if appropriate, HER2/NEU-targeted therapy.

Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Through end of benefit year.

#### Drugs LORBRENA ORAL TABLET 100 MG, 25 MG

# **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Documentation of anaplastic lymphoma kinase (ALK)-positive metastatic non-small cell lung cancer (NSCLC) whose disease has progressed on one of the following: 1. crizotinib and at least one other ALK inhibitor for metastatic disease: or 2. alectinib as the first ALK inhibitor therapy for metastatic disease: or 3. ceritinib as the first ALK inhibitor therapy for metastatic disease.

# Age Restriction

Prescriber Restriction Oncology

**Coverage Duration** 3 months

Drugs LUPKYNIS

**Covered Uses** 

**Exclusion Criteria** 

#### **Required Medical Information**

Diagnosis of Lupus nephritis, Active, in combination with mycophenolate mofetil and corticosteroids

Age Restriction 18 years of age and older

**Prescriber Restriction** 

**Coverage Duration** 1 year

# **Other Criteria**

Establish an accurate baseline estimated GFR (eGFR) and blood pressure (BP) prior to initiation. Do not initiate treatment in patient with BP greater than 165/105 mm Hg or with hypertensive emergency. Use not recommended with baseline eGFR of 45 mL/min/1.73m(2)

# Drugs

# LUPRON DEPOT, LUPRON DEPOT (3 MONTH), LUPRON DEPOT (4 MONTH), LUPRON DEPOT (6 MONTH), LUPRON DEPOT-PED, LUPRON DEPOT-PED (3 MONTH)

#### **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

Pregnancy in patients with child-bearing potential. Breastfeeding. Undiagnosed abnormal vaginal bleeding.

# **Required Medical Information**

Diagnosis of one of the following: A) Advanced or metastatic prostate cancer (7.5 mg 1-month, 22.5 mg 3-month, 30 mg 4-month, & 45 mg 6-month depots only), or B) Endometriosis (3.75 mg 1-month & 11.25 mg 3-month depots only) AND 1. For initial authorization, patient has had an inadequate pain control response or has an intolerance or contraindication to one of the following: Danazol or combination [estrogen/progesterone] oral contraceptives or progestins, or 2. For retreatment course, patient is experiencing recurrence of symptoms after an initial course of therapy with leuprolide acetate and norethindrone acetate 5 mg daily will be co-administered, or C) Anemia due to uterine Leiomyomata (Fibroids) (3.75 mg 1-month &11.25 mg 3-month depots only) AND patient is preoperative AND has tried and had an inadequate response to monotherapy with iron, or D) Central precocious puberty (Lupron Depot-Ped) AND submission of pubertal gonadal sex steroid levels (testosterone greater than 30 ng/dL, estradiol greater than 20 pg/mL AND a pubertal LH increase upon native GnRH stimulation AND pelvic ultrasound assessment (girls) is required for approval along with notes indicating premature development of secondary sexual characteristics at or before the age of 8 yrs in girls and 9 yrs in boys and significant advancement of bone age and/or a poor adult height prediction AND other causes of sexual precocity must be excluded.

# **Age Restriction**

# **Prescriber Restriction**

Oncologist, Endocrinologist, or Gynecologist to prescribe

# **Coverage Duration**

Plan year

#### **Other Criteria**

For endometriosis and uterine fibroids, patient will be using nonhormonal contraception during and for 12 weeks after therapy.

Drugs LYNPARZA

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

1. Documentation of deleterious germline BRCA mutated ovarian cancer AND 2. Documentation of at least 3 prior chemotherapy regimens that have been ineffective or not tolerated AND 3. Lynparza will be used as monotherapy.

# **Age Restriction**

**Prescriber Restriction** 

**Coverage Duration** Through end of benefit year

#### Drugs MAKENA (PF)

# **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

#### **Required Medical Information**

Clinical documentation of singleton pregnancy (i.e. one fetus) AND a history of singleton spontaneous preterm birth defined as delivery prior to 37 weeks gestation AND the pregnancy is between 16 weeks, 0 days and 20 weeks, 6 days gestation AND the requested dose and frequency is in accordance with FDA-approved labeling.

# Age Restriction

16 years of age or older

# **Prescriber Restriction**

**Coverage Duration** coverage is provided until week 37 (through 36 weeks, 6 days) of gestation

Drugs

MAVENCLAD (10 TABLET PACK), MAVENCLAD (4 TABLET PACK), MAVENCLAD (5 TABLET PACK), MAVENCLAD (6 TABLET PACK), MAVENCLAD (7 TABLET PACK), MAVENCLAD (8 TABLET PACK), MAVENCLAD (9 TABLET PACK)

#### **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D.

#### **Exclusion Criteria**

Individual with current malignancy OR human immunodeficiency virus (HIV) infection OR an active chronic infection (e.g., hepatitis or tuberculosis)

#### **Required Medical Information**

Documentation of diagnosis of relapsing multiple sclerosis, including relapsing-remitting disease or active secondary progressive disease AND the patient has had a trial and inadequate response or intolerance to at least one alternative drug indicated for the treatment of multiple sclerosis.

**Age Restriction** 

Prescriber Restriction Coverage Duration

1 year

Other Criteria Will only be approved for 2 treatment cycles.

#### Drugs MAVYRET ORAL TABLET

# **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

For retreatment, patient was non-adherent to initial regimen as evidenced by medical record and/or pharmacy claims OR patient continues to engage in high risk behavior and experienced reinfection secondary to high risk behavior.

# **Required Medical Information**

Provider must submit medical records documenting the following: 1) medical diagnosis of Chronic Hepatitis C with labs documenting genotype and subtype, AND 2) medical records documenting viral load taken within 6 months of beginning therapy, AND 3) documentation of previous HCV therapies to confirm appropriate duration of treatment. Authorization for retreatment requires the following: 1)Evidence of failure to achieve a sustained virologic response (SVR) or lack of efficacy during treatment (polymerase chain reaction (PCR) assay, 12 or more weeks after completing treatment or a 10-fold increase of viral load at week 6 of treatment) OR evidence of adverse event that required therapy discontinuation (Laboratory results and/or clinical presentation), AND 2) Member was adherent to previous therapy as evidenced by pharmacy claims, AND 3) Submission of psychological support/treatment for a minimum of six months for substance abuse related failure (i.e. NA, AA), AND 4)Patient has abstained from the use of illicit drugs and alcohol for a minimum of 3 months as evidenced by negative urine or blood confirmation tests, collected monthly for the past 90 days prior to initiation of therapy.

# **Age Restriction**

#### **Prescriber Restriction**

Prescribed by, a gastroenterologist, hepatologist, or infectious disease physician.

#### **Coverage Duration**

based on the AASLD treatment guidelines

#### **Other Criteria**

Criteria and coverage durations will be applied consistent with current AASLD/IDSA guidance.

#### Drugs MAYZENT ORAL TABLET 0.25 MG, 2 MG, MAYZENT STARTER(FOR 2MG MAINT)

# **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D. Diagnosis of relapsing forms of multiple sclerosis and must have tried and failed one generic alternative agent such as dimethyl fumarate or glatiramer acetate.

**Exclusion Criteria** 

**Required Medical Information** 

Age Restriction

**Prescriber Restriction** 

**Coverage Duration** 1 year

#### MEKINIST

#### Drugs MEKINIST ORAL TABLET 0.5 MG, 2 MG

# **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Diagnosis of unresectable or metastatic melanoma, positive BRAF V600E or V600K mutation as detected by an FDAapproved test (THxID-BRAF Kit) or Clinical Laboratory Improvement Amendments (CLIA)-approved facility, and the patient has not received prior BRAF-inhibitor therapy.

Age Restriction 18 years or older

Prescriber Restriction Oncologist

**Coverage Duration** 3 months

Drugs MEKTOVI

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Documentation of unresectable or metastatic melanoma with a BRAF V600E or V600K mutation, as detected by an FDAapproved test and that Mektovi will be used in combination with encorafenib.

# Age Restriction

Prescriber Restriction Oncology

**Coverage Duration** 3 months

#### Drugs MENEST

# **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

Any pertinent clinical situation as defined by the product label that could affect patient safety and/or therapeutic efficacy (i.e. contraindications, warnings, precautions, adverse effects, renal or hepatic function, drug interactions, lab values, required prior or concomitant therapy, inappropriate dosing and/or duration, etc).

#### **Required Medical Information**

Clinical documentation of FDA approved indication for treatment.

Age Restriction 18 years and older

**Prescriber Restriction** 

Coverage Duration Plan year

# Other Criteria

For treatment of vaginal atrophy - must have documented failure, intolerance or contraindication to at least 1 formulary vaginal estrogen. (Estrace cream, Premarin cream, Vagifem tab)

# Miglustat

Drugs miglustat

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Documentation of mild to moderate type 1 Gaucher disease AND patient is symptomatic (i.e. radiologic evidence of skeletal disease, platelet count less than 60,000 microL, liver greater than 2.5 times normal size, spleen greater than 15 times normal size) AND enzyme replacement therapy is not a therapeutic option (e.g. due to allergy, hypersensitivity, or poor venous access).

Age Restriction 18 years of age or older

**Prescriber Restriction** 

**Coverage Duration** 3 Months

**Drugs** modafinil

**Covered Uses** All FDA-Approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

**Required Medical Information** Documentation of diagnosis and prior therapies used

# **Age Restriction**

#### **Prescriber Restriction**

Idiopathic hypersomnia-- approve if the diagnosis is confirmed by a sleep specialist physician or at an institution that specializes in sleep disorders (i.e., sleep center)

#### **Coverage Duration** Plan year

Drugs MONJUVI

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Patient has a diagnosis of diffuse large B-cell lymphoma (DLBCL) not otherwise specified, including DLBCL arising from low grade lymphoma (excluding primary refractory AND 'double or triple hit' disease) AND Patient has at least one measurable lesion AND Patient's disease is relapsed or refractory AND Therapy will be initiated in combination with lenalidomide AND Patient is NOT eligible for autologous stem cell transplant AND Patient is ineligible for intensive therapy (i.e., high-dose chemotherapy (HDC) and autologous stem cell transplantation (ASCT))

Age Restriction Patient is 18 years or older

**Prescriber Restriction** Prescribed by oncologist

**Coverage Duration** 3 months

#### Drugs MOVANTIK

#### **Covered Uses**

All FDA-approved indications not otherwise excluded from Health Plan

#### **Exclusion Criteria**

Known or suspected gastrointestinal obstruction and at increased risk of recurrent obstruction, due to the potential for gastrointestinal perforation. Concomitantly taking strong CYP3A4 inhibitors (e.g., clarithromycin, ketoconazole)

#### **Required Medical Information**

Documentation of diagnosis and treatment history.

**Age Restriction** 

**Prescriber Restriction** 

Coverage Duration Plan year

# **Other Criteria**

Approve if member has been taking an opioid for at least 4 weeks and has tried lifestyle changes (e.g. maintaining a diet rich in fiber and/or fiber supplementation along with adequate fluid intake) and has tried a bowel regimen of an osmostic laxative (e.g. PEG 3350) or a stimulant laxative (e.g. bisacodyl) with or without a stool softener (e.g. docusate).

Drugs MULPLETA

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Documentation that the medication will be used for the treatment of thrombocytopenia in adult patients with chronic liver disease who are scheduled to undergo a procedure and had a platelet count less than 50 × 10^9/L

Age Restriction

**Prescriber Restriction** 

**Coverage Duration** 3 Months

Drugs NAGLAZYME

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

**Required Medical Information** 

# **Age Restriction**

**Prescriber Restriction** 

**Coverage Duration** Plan year

Drugs NERLYNX

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

NERLYNX is indicated for the extended adjuvant treatment of adult patients with early stage HER2overexpressed/amplified breast cancer, to follow adjuvant trastuzumab based therapy.

# Age Restriction

Prescriber Restriction Oncologist

**Coverage Duration** Through the end of benefit year

# Drugs NEULASTA, NEULASTA ONPRO

# **Covered Uses**

All FDA-approved indications not otherwise excluded from Health Plan.

# **Exclusion Criteria**

**Required Medical Information** Documentation of diagnosis.

# **Age Restriction**

**Prescriber Restriction** 

**Coverage Duration** 3 months and is renewable in situations where it continues to provide clinical benefit

Drugs NEUPOGEN

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

**Exclusion Criteria** 

**Required Medical Information** Clinical documentation of FDA approved indication for treatment.

Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Varies by indication.

**Drugs** nevirapine

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

**Required Medical Information** Clinical documentation of FDA approved indication for treatment.

# Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Plan year

#### Drugs NEXAVAR

# **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

Combination use with other tyrosine kinase inhibitors such as sorafenib, sunitinib. Squamous cell lung cancer being treated with carboplatin and paclitaxel.

#### **Required Medical Information**

Diagnosis of one of the following: A) Advanced renal cell carcinoma, or B) Locally recurrent or metastatic, progressive, differentiated thyroid carcinoma refractory to radioactive iodine treatment, or C) Unresectable hepatocellular carcinoma.

#### **Age Restriction**

Patient must be at least 18 years old or older.

#### **Prescriber Restriction**

#### **Coverage Duration** Initial: 3 months, Renewal: through end of benefit year w/ stable disease

Drugs NEXLETOL

**Covered Uses** 

**Exclusion Criteria** 

#### **Required Medical Information**

Clinical documentation (e.g., chart notes, laboratory values) required for initial authorization includes: 1. Documentation of one of the following diagnoses: A) Primary Hyperlipidemia (Including Heterozygous Familial Hypercholesterolemia), OR B) Atherosclerotic cardiovascular disease (ASCVD), AND 2. Documentation of any one of the following: a) Patient has been receiving statin therapy at a maximally tolerated dose and (maximally tolerated dose may include no statin therapy) OR b) Patient has a documented labeled contraindication to all statins OR c) Patient has experienced rhabdomyolysis, AND 3. Patient has tried and failed ezetimibe. Clinical documentation required for reauthorization includes: 1. Physician attestation of medication's efficacy.

**Age Restriction** 

18 years or older

#### **Prescriber Restriction**

**Coverage Duration** Through end of benefit year

#### **Other Criteria**

Other Criteria: Primary Hyperlipidemia (Including Heterozygous Familial Hypercholesterolemia): The patient has a lowdensity lipoprotein cholesterol (LDL-C level) greater than or equal to 100 mg/dL. C) ASCVD: The patient has a lowdensity lipoprotein cholesterol (LDL-C) greater than or equal to 70 mg/dL (after treatment with antihyperlipidemic agents. ASCVD diagnosis is confirmed by one of the following: acute coronary syndromes, history of myocardial infarction, stable or unstable angina, coronary or other arterial revascularization, stroke, transient ischemic attack, peripheral arterial disease presumed to be of atherosclerotic origin. Drugs NEXLIZET

**Covered Uses** 

**Exclusion Criteria** 

#### **Required Medical Information**

Clinical documentation (e.g., chart notes, laboratory values) required for initial authorization includes: 1. Documentation of one of the following diagnoses: A) Heterozygous familial hypercholesterolemia, OR B) Atherosclerotic cardiovascular disease (ASCVD) established, AND 2. Documentation of any one of the following: a) Patient has been receiving statin therapy at a maximally tolerated dose (maximally tolerated dose may include no statin therapy), OR b) Patient has a documented labeled contraindication to all statins, OR c) Patient has experienced rhabdomyolysis. Clinical documentation required for reauthorization includes: 1. Physician attestation of medication's efficacy.

# Age Restriction

18 years and older.

#### **Prescriber Restriction**

# **Coverage Duration**

Through end of benefit year

# **Other Criteria**

Other Criteria: Primary Hyperlipidemia (Including Heterozygous Familial Hypercholesterolemia): The patient has a lowdensity lipoprotein cholesterol (LDL-C level) greater than or equal to 100 mg/dL. C) ASCVD: The patient has a lowdensity lipoprotein cholesterol (LDL-C) greater than or equal to 70 mg/dL (after treatment with antihyperlipidemic agents. ASCVD diagnosis is confirmed by one of the following: acute coronary syndromes, history of myocardial infarction, stable or unstable angina, coronary or other arterial revascularization, stroke, transient ischemic attack, peripheral arterial disease presumed to be of atherosclerotic origin. Drugs NINLARO

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Diagnosis of multiple myeloma AND have received at least one prior therapy AND medication will be used in combination with dexamethasone and lenalidomide.

# Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Through end of benefit year.

# Drugs NORPACE CR

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

# **Required Medical Information**

Documentation of diagnosis and treatment history including reason why disopyramide IR cannot be used.

# **Age Restriction**

**Prescriber Restriction** 

**Coverage Duration** Plan year

Drugs

droxidopa

Covered Uses

All FDA-approved indications not otherwise excluded from Health Plan.

#### **Exclusion Criteria**

# **Required Medical Information**

Age Restriction 18 years of age or older

# **Prescriber Restriction**

**Coverage Duration** Plan year

# **Other Criteria**

For renewal, Patient does not have persistent or sustained supine hypertension (SBP more than 180 mmHg or DBP more than 110 mmHg), Patient does not have persistent or sustained standing or sitting hypertension (SBP more than 180 mmHg or DBP more than 110 mmHg), and Patient had improvement in symptoms of NOH. Sustained mean elevated blood pressure that persists for longer than 5 minutes after change in position. Persistent means elevated BP that occurs on more than one occasion on separate physician office visits

# Drugs NORVIR ORAL SOLUTION

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

**Required Medical Information** Clinical documentation of FDA approved indication for treatment.

Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Plan year

# Drugs NOXAFIL

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

# **Required Medical Information**

Clinical documentation of FDA approved indication for treatment.

**Age Restriction** 13 years and older.

# **Prescriber Restriction**

# **Coverage Duration**

Per treatment OR up to through plan year.

# **Other Criteria**

Fluconazole preferred for candida. Voriconazole preferred for aspergillus.

Drugs NUBEQA

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

**Exclusion Criteria** 

**Required Medical Information** Documenation of non-metastatic castration resistant prostate cancer (nmCRPC).

Age Restriction

Prescriber Restriction Oncologist

**Coverage Duration** 3 months

#### Drugs NUCALA SUBCUTANEOUS AUTO-INJECTOR

# **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Documentation that either A) patient has asthma with an eosinophilic phenotype defined as blood eosinophils greater than or equal to 300 cells/µL within previous 12 months or greater than or equal to 150 cells/µL within 6 weeks of dosing AND medication will be used in combination with a corticosteroid inhaler and long acting beta2-agonist AND Patient must have experienced two or more exacerbations in the previous year OR require daily oral corticosteroids OR B)patient has eosinophilic granulomatosis with polyangiitis (EGPA)AND documented trial and failure of or contraindication to treatment with at least one immunosuppressants (azathioprine, cyclophosphamide, or methotrexate).

Age Restriction

**Prescriber Restriction** 

**Coverage Duration** 6 months

# Drugs NUCYNTA, NUCYNTA ER

**Covered Uses** 

All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

**Required Medical Information** Documentation of diagnosis and treatment history.

# Age Restriction 18 years and older.

# **Prescriber Restriction**

**Coverage Duration** Plan year

**Other Criteria** Documented failure to tramadol or tramadol extended-release. Drugs NULOJIX

#### **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

Transplant recipients who are Epstein-Barr virus (EBV) seronegative or with unknown EBV serostatus.

#### **Required Medical Information**

For prophylaxis of organ rejection in adults receiving kidney transplant, in combination with basiliximab induction, mycophenolate mofetil, and corticosteroids, AND documentation of patient's EBV serostatus.

# Age Restriction 18 years of age or older

# **Prescriber Restriction**

**Coverage Duration** Plan year

Drugs NURTEC ODT

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Documentation that the medication will be used for the acute treatment of migraine with or without aura in adults AND member has tried and failed two triptan therapies OR Documentation that the medication will be used for the prevention of migraine with or without aura in adults AND member has tried and failed two preventative medication alternatives

# **Age Restriction**

**Prescriber Restriction** 

# Coverage Duration

Through end of benefit year

#### **Other Criteria**

Reauthorization requires documentation of medication efficacy.

#### Drugs

octreotide acetate injection solution

#### **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

#### **Required Medical Information**

Diagnosis of one of the following: A) Acromegaly and patient had an inadequate response or cannot be treated with surgical resection, pituitary irradiation, and/or bromocriptine mesylate at maximally tolerated doses, or B) Metastatic carcinoid tumor requiring symptomatic treatment of severe diarrhea and flushing episodes, or C) Vasoactive intestinal peptide tumor requiring treatment of profuse watery diarrhea. Acromegaly: Documentation of inadequate response to surgery and/or radiotherapy, or documentation that patient is not a candidate for surgery and/or radiotherapy. Reauthorization will require statement indicating growth hormone (GH) levels are stabilized at less than 5.0ng/mL and IGF-1 levels are normalized (male less than 1.9U/mL or female less than 2.2 U/mL) as matched by age and gender, or the patient has a documented clinical response defined by a reduction of tumor mass, a reduction in the signs and symptoms of acromegaly, or an improvement in significant comorbidities.

#### Age Restriction

**Prescriber Restriction** 

Coverage Duration Plan year

#### **Other Criteria**

For renewal of acromegaly, IGF-1 level has normalized or improved. For renewal of metastatic carcinoid tumor, patient has improvement in diarrhea and flushing episodes. For renewal of vasoactive intestinal peptide tumor, improvement in diarrhea episodes.

Drugs ODEFSEY

**Covered Uses** All FDA-approved indications not otherwise excluded from Health Plan

**Exclusion Criteria** 

**Required Medical Information** 

Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Plan year

Drugs ODOMZO

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Documented locally advanced basal cell carcinoma (BCC) that has recurred following surgery or radiation therapy, or the member is a not candidate for surgery or radiation therapy.

# Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Through end of benefit year.

#### Drugs OMNIPOD CLASSIC PDM KIT(GEN 3), OMNIPOD CLASSIC PODS (GEN 3), OMNIPOD DASH PDM KIT (GEN 4), OMNIPOD DASH PODS (GEN 4)

## **Covered Uses**

**Exclusion Criteria** 

## **Required Medical Information**

Initial Therapy 1)Diagnosis of Type 1 Diabetes2)Member has previously been on a maintenance program involving at least three injections of insulin per day and frequent self-adjustments of insulin dosage or current use of an insulin pump3)Member performs glucose self-testing at least three times per day4)History of suboptimal blood sugar control despite appropriate management5)Member or caregiver has completed a physician-directed comprehensive diabetes management programContinued Therapy1)There is documented evidence of improvement in control of diabetes (specific to baseline status of disease for individual members)

Age Restriction None

Prescriber Restriction Endocrinologist

**Coverage Duration** 12 months

### **Other Criteria**

If request is for more than 10 pods per 30 days additional documentation is required to provide clinical rationale for higher quantity

Drugs ONUREG

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Documented diagnosis of Acute myeloid leukemia AND achieved first complete remission (CR) or complete remission with incomplete blood count recovery (CRi) following intensive induction chemotherapy AND are not able to complete intensive curative therapy AND Have tried and failed or has contraindication to all other NCCN recommended therapies (Azacitadine SQ or IV, Decitabine, Low-dose cytarabine, Venclexta) AND Baseline CBC tests be performed prior to initiation.

**Age Restriction** 

**Prescriber Restriction** Oncologist or hematologist

**Coverage Duration** 3 months

#### Drugs ORFADIN ORAL CAPSULE 10 MG, 2 MG, 5 MG

# **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

## **Required Medical Information**

Medical statement of diagnosis of hereditary tyrosinemia type 1 (HT-1) AND current patient weight as dose must be within FDA approved dosing range: maximum dosage for all patients is 2 mg/kg/day. When initiating therapy, Serum tyrosine should be below 500 mmol/L to avoid toxic effects, and urinary succinylacetone levels should be undetectable.

### **Age Restriction**

**Prescriber Restriction** 

**Coverage Duration** 3 Months

#### Drugs ORIAHNN

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

Presence of any contraindications as defined by the FDA.

# **Required Medical Information**

Documentation that the medication will be used for an FDA approved indication and Trial of two previous therapies (estrogen/progestin contraceptives, Progestin IUD, tranexamic acid).

Age Restriction 18 years of age or older

**Prescriber Restriction** physician specializing in patient's diagnosis or is in consultation with a gynecologist

**Coverage Duration** 12 months

Other Criteria Max duration of therapy for 2 years

#### Drugs ORILISSA ORAL TABLET 150 MG, 200 MG

# **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

presence of medication contraindication as defined by the FDA.

# **Required Medical Information**

Documentation that the medication will be used for an FDA approved indication AND trial of NSAIDs, estrogen/progestin unless contraindicated ANDtrial of Eligard unless contraindication.

Age Restriction 18 years of age or older

# **Prescriber Restriction**

physician specializing in patient's diagnosis or is in consultation with a gynecologist

**Coverage Duration** 12 months

Other Criteria

Treatment not to exceed two years in duration

#### **ORKAMBI** (lumacaftor-ivacaftor)

Drugs ORKAMBI

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

#### **Exclusion Criteria** Use in combination with Kalydeco

# **Required Medical Information**

The patient is positive for the F508del mutation on both alleles of the CFTR gene.

Age Restriction 6 years of age or older

**Prescriber Restriction** 

**Coverage Duration** Through end of benefit year.

# Drugs OTEZLA, OTEZLA STARTER

# Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

#### **Required Medical Information** Documented patient is free of any clinically important active infections.

# Age Restriction

**Prescriber Restriction** 

**Coverage Duration** 12 months

#### Drugs

oxandrolone oral tablet 10 mg, 2.5 mg

#### **Covered Uses**

All medically accepted indications not otherwise excluded from Health Plan.

#### **Exclusion Criteria**

Pregnancy Category X

# **Required Medical Information**

Statement indicating use to promote weight gain after weight loss following extensive surgery, chronic infections, or severe trauma, and in some patients who without definite pathophysiologic reasons fail to gain or to maintain normal weight, and to offset the protein catabolism associated with prolonged administration of corticosteroids. Statement indicating use for orphan drug indication, short stature associated with Turner syndrome, constitutional delay of growth and puberty, moderate or severe acute alcoholic hepatitis, Duchenne and Becker muscular dystrophy. Initial Therapy for AIDS Wasting: Diagnosis of AIDS wasting/cachexia. For treatment of anorexia associated with weight loss in patients with HIV: 1. Patient is receiving AIDS anti-retroviral therapy AND 2. experienced as least a. 7.5% unintentional weight loss over 6 months b. 10% unintentional weight loss over 12 months c. 5% body cell mass (BCM) loss within 6 months d. BMI less than 20 kg/m2 e. BCM less than 35% male (less than 23% female) and a BMI less than27 kg/m2 AND 3. documentation of trial and failure, contraindication, or intolerance to megestrol at doses up to 800mg daily.

#### **Age Restriction**

**Prescriber Restriction** 

**Coverage Duration** HIV Wasting: 3 months. All other indications: Through the Benefit Year

Drugs OXERVATE

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Member has a diagnosis (documented in chart notes) of stage 2 (recurrent/persistent epithelial defect) or stage 3 (corneal ulcer) neurotrophic keratitis in the affected eye(s) AND Member is refractory to at least ONE conventional non-surgical treatment for neurotrophic keratitis (e.g. preservative-free artificial tears, topical antibiotic eyedrops, therapeutic contact lenses, etc.)

Age Restriction Member is 2 years of age or older

#### **Prescriber Restriction**

The medication is prescribed by an ophthalmologist

**Coverage Duration** 8 weeks

Drugs OXLUMO

**Covered Uses** 

All FDA-approved indications not otherwise excluded from Part D.

## **Exclusion Criteria**

## **Required Medical Information**

Documented diagnosis of PH1 by, or in consultation with, a specialist (e.g., geneticist, nephrologist, urologist) with expertise in the diagnosis of PH1) AND confirmation of the PH1 diagnosis based on both metabolic (Increased urinary oxalate excretion greater than 1 mmol/1.73 m2 per day [90 mg/1.73 m2 per day OR increased urinary oxalate: creatinine ratio relative to normative values for age) OR increased plasma oxalate and glyoxylate concentrations) AND Genetic testing confirming a mutation in the alanine:glyoxylate aminotransferase (AGT or AGXT) gene AND patient has not received a liver transplant AND Oxlumo dosing is in accordance with the United States Food and Drug Administration approved labeling

#### **Age Restriction**

#### **Prescriber Restriction**

Oxlumo is prescribed by, or in consultation with, a specialist (e.g., geneticist, nephrologist, urologist) with expertise in the treatment of PH1

**Coverage Duration** 6 months

Drugs PALYNZIQ

# **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D.

#### **Exclusion Criteria**

Not to be used in combination with sapropterin dihydrochloride (Kuvan)

### **Required Medical Information**

Documented Diagnosis of Phenylketonuria (PKU), AND blood phenylalanine concentration greater than 600 micromol/L, AND physician agrees to assess patient tolerability, blood phenylalanine concentration, dietary protein and phenylalanine intake throughout treatment, AND prescriber and patient must be enrolled with the Palynziq REMS Program.

Age Restriction 18 years of age and older

**Prescriber Restriction** 

**Coverage Duration** 3 Months

#### PAMIDRONATE

Drugs

pamidronate

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

#### **Required Medical Information**

Clinical documentation of FDA approved indication for treatment and treatment history. For treatment of hypercalcemia of malignancy, documentation of corrected total serum calcium greater than or equal to 12 mg/dL. For treament of bone metastases, diagnosis of breast cancer or multiple myeloma. For Paget's disease, must have symptomatic form of disease.

Age Restriction 18 years and older

#### **Prescriber Restriction**

**Coverage Duration** Plan year

#### **Other Criteria**

For Paget's disease, must have documented failure, intolerance or contraindication to oral agents: alendronate or risedronate.

Drugs PANRETIN

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

# **Required Medical Information**

Age Restriction 18 years and older

Prescriber Restriction Oncologist or HIV specialist

**Coverage Duration** Plan year

#### paricalcitol (ZEMPLAR)

Drugs

paricalcitol oral

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

### **Exclusion Criteria**

**Required Medical Information** Documentation of diagnosis and treatment history.

Age Restriction 18 years and older.

Prescriber Restriction

Coverage Duration Plan year

**Other Criteria** Documented failure or intolerance to calcitriol.

# Penicillamine capsule

Drugs penicillamine oral capsule, penicillamine oral tablet

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

**Required Medical Information** 

#### **Age Restriction**

**Prescriber Restriction** 

**Coverage Duration** 3 months

Drugs PENTAM

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

**Required Medical Information** Clinical documentation of FDA approved indication for treatment.

Age Restriction 4 months and older

**Prescriber Restriction** 

Coverage Duration Per treatment

Drugs PERSERIS

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

**Required Medical Information** Documented trial of any of the two following drugs: aripiprazole, clozapine, olanzapine, risperidone, quetiapine, ziprasidone

**Age Restriction** 

**Prescriber Restriction** 

**Coverage Duration** Through the benefit year

# Drugs PIQRAY ORAL TABLET 200 MG/DAY (200 MG X 1), 250 MG/DAY (200 MG X1-50 MG X1), 300 MG/DAY (150 MG X 2)

### **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D.

#### **Exclusion Criteria**

#### **Required Medical Information**

Documentation that the patient is a postmenopausal female or a male AND has advanced or metastatic breast cancer AND has HR-positive disease AND has HER2-negative disease AND has PIK3CA-mutated breast cancer as detected by a FDA approved test AND has progressed on or after at least one prior endocrine-based regimen AND the medication will be used in combination with fulvestrant.

## Age Restriction

Prescriber Restriction Oncologist

**Coverage Duration** 3 Months

# Drugs PLEGRIDY SUBCUTANEOUS

#### **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Diagnosis of relapsing forms of multiple sclerosis and must have tried and failed one generic alternative agent such as dimethyl fumarate or glatiramer acetate

Age Restriction Age 18 years and older

Prescriber Restriction Neurologist

**Coverage Duration** 6 months

Drugs POMALYST

**Covered Uses** All FDA-approved indications not otherwise excluded from Health Plan.

# **Exclusion Criteria**

# **Required Medical Information**

Patient has a diagnosis of multiple myleoma and the patient has received two prior therapies, including Revlimid and Velcade unless the patient has a contraindication or intolerance to Revlimid or Velcade and the patient has demonstrated disease progression on or within 60 days of completion of last therapy.

#### **Age Restriction**

**Prescriber Restriction** 

**Coverage Duration** 3 Months

#### **Other Criteria**

Prescriber, pharmacist, and patient must be enrolled in the Pomalyst REMS program.

# Drugs POTELIGEO

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

**Required Medical Information** Documentation of relapsed or refractory mycosis fungoides (MF) or Sézary syndrome (SS) after at least one prior systemic therapy.

Age Restriction 18 years or older

## **Prescriber Restriction**

**Coverage Duration** 3 Months

#### Drugs

pramipexole oral tablet extended release 24 hr 0.375 mg, 0.75 mg, 1.5 mg, 3 mg, 4.5 mg

# Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

## **Exclusion Criteria**

# Required Medical Information

Documented diagnosis and treatment history.

#### Age Restriction 18 years and older

#### **Prescriber Restriction**

**Coverage Duration** Plan year

Other Criteria Failure to pramipexole IR Drugs PREZCOBIX

**Covered Uses** All FDA-approved indications not otherwise excluded from Health Plan

**Exclusion Criteria** 

**Required Medical Information** 

Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Plan year

# Drugs PREZISTA ORAL SUSPENSION, PREZISTA ORAL TABLET 150 MG, 600 MG, 75 MG, 800 MG

#### **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

**Exclusion Criteria** 

## **Required Medical Information**

Clinical documentation of FDA approved indication for treatment.

**Age Restriction** 

**Prescriber Restriction** 

Coverage Duration Plan year

# Drugs

# PROCRIT INJECTION SOLUTION 10,000 UNIT/ML, 2,000 UNIT/ML, 20,000 UNIT/ML, 3,000 UNIT/ML, 4,000 UNIT/ML, 40,000 UNIT/ML

#### **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

Uncontrolled hypertension. Pure red cell aplasia that begins after ESA treatment.

#### **Required Medical Information**

Pre-treatment hemoglobin level less than 10 g/dL AND Patient has adequate iron stores prior to initiation of therapy defined as ferritin more than 100 mcg/L or serum transferrin saturation greater than 20% AND other causes of anemia such as iron deficiency, folate deficiency or B12 deficiency, hemolysis, gastrointestinal bleeding, other active or occult bleeding, or underlying hematologic disease (such as sickle cell anemia, thalassemia, and porphyria) have been ruled out AND Diagnosis of one of the following: A) Anemia due to chronic kidney disease (CKD) with or without hemodialysis, OR B) Anemia in patients with non-myeloid malignancies where anemia is due to the effect of concomitant myelosuppressive chemotherapy and two additional months of chemotherapy is anticipated, C) Treatment of anemic in a patient at high risk for perioperative blood loss from elective, noncardiac, nonvascular surgery to reduce the need for allogeneic blood transfusion, D) Anemia in zidovudine-treated HIV infection with serum erythropoietin levels 500 mUnits/mL or less and zidovudine doses 4,200 mg/week or less.

# **Age Restriction**

#### **Prescriber Restriction**

CKD - prescribed by a nephrologist or hematologist. Non-myeloid malignancies - prescribed by an oncologist/hematologist. Surgery - Prescribed by a surgeon. HIV - Prescribed by an infectious disease specialist.

#### **Coverage Duration**

Initial: 3 months. Renewal: CKD-12 months, Non-myeloid cancers, HIV-4 months. Surgery-3 months

#### **Other Criteria**

For renewal of CKD (dialysis patients): Hb less than 11 g/dL or physician will decrease or interrupt dose. For renewal of CKD (non-dialysis patients): Hb less than 10 g/dL or physician will decrease or interrupt dose. For renewal of non-myeloid malignancies: Concurrent myelosuppressive chemotherapy and Hb is 12g/dL or less and there is measurable response after eight weeks (defined as an increase in Hb 1 g/dL or more or a reduction in red blood cell transfusion requirements). For renewal of zidovudine-treated HIV, Hb is 12g/dL or less AND Zidovudine dose remains 4,200 mg/week or less and there is a measurable response after eight weeks (defined as an increase in Hb or a reduction in RBC transfusion requirements or documented dose escalation [up to max of 300 units/kg/dose]).

#### Drugs PROMACTA ORAL POWDER IN PACKET 12.5 MG, PROMACTA ORAL TABLET

# **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

# **Required Medical Information**

Diagnosis of one of the following: A) Relapsed/refractory chronic immune (idiopathic) thrombocytopenic purpura (ITP) for greater than 6 months AND Baseline platelet count is less than 50,000/mcL AND Degree of thrombocytopenia and clinical condition increase the risk of bleeding AND Patient had an insufficient response, intolerance, contraindication to corticosteroids or immune globulin or inadequate response or contraindication to splenectomy, or B) Chronic hepatitis C and patient has thrombocytopenia defined as platelets less than 90,000/mcL for initiation (pre-treatment) of interferon therapy, or C) Severe aplastic anemia and patient has insufficient response to immunosupressive therapy.

Age Restriction

**Prescriber Restriction** 

**Coverage Duration** 3 Months

# **PROTOPIC** (tacrolimus topical)

Drugs

tacrolimus topical

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

# **Required Medical Information** Clinical documentation of FDA approved indication for treatment.

Age Restriction 16 years and older (0.1%) or 2 years and older (0.03%)

# **Prescriber Restriction**

**Coverage Duration** 8 weeks

Drugs PULMOZYME

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

**Exclusion Criteria** 

**Required Medical Information** Clinical documentation of Cystic Fibrosis diagnosis

Age Restriction

**Prescriber Restriction** 

Coverage Duration Plan year

# **Other Criteria**

Pulmozyme should be used in conjunction with standard therapies for CF. For renewal, Patient is benefiting from treatment (i.e. improvement in lung function [FEV1], decreased number of pulmonary exacerbations).

Drugs REGRANEX

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

**Required Medical Information** Documentation of diagnosis

Age Restriction 16 years and older

**Prescriber Restriction** 

**Coverage Duration** Plan year

# Drugs RELENZA DISKHALER

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

**Required Medical Information** Clinical documentation of FDA approved indication for treatment.

# Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Per treatment (up to 28 days)

#### Drugs REMODULIN

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

# **Required Medical Information** Clinical documentation of FDA approved indication for treatment.

Age Restriction 18 years and older.

**Prescriber Restriction** 

**Coverage Duration** 3 months

#### Drugs REPATHA PUSHTRONEX, REPATHA SURECLICK, REPATHA SYRINGE

# **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D. All FDA-approved indications not otherwise excluded by Health Plan.

### **Exclusion Criteria**

Concurrent use of Repatha with Praluent (alirocumab injection for SC use), Juxtapid (lomitapide capsules) or Kynamro (mipomersen injection). The efficacy and safety of Praluent, Juxtapid and Kynamro in combination with Repatha have not been established.

#### **Required Medical Information**

For "initial" authorization: 1. One of the following diagnoses: A) Homozygous familial hypercholesterolemia (HoFH), B) Heterozygous familial hypercholesterolemia (HeFH), OR C) Atherosclerotic cardiovascular disease (ASCVD), AND 2. Documentation of any one of the following: a) Patient has been receiving at least 12 consecutive weeks of high intensity statin therapy and will continue to receive high-intensity statin at maximally tolerated dose, b) Patient is unable to tolerate high-intensity statin, c) Patient has been receiving at least 12 consecutive weeks of moderate-intensity statin therapy and will continue to receive a moderate-intensity statin d) Patient has a documented labeled contraindication to all statins, e)Patient has experienced rhabdomyolysis or muscle symptoms with statin treatment, f)Patient has undergone a trial of statin rechallenge with another low-intensity statin with documented reappearance of muscle symptoms (only 2 trials of a statin can be required prior to the approval of a PCSK9). Clinical documentation required for "reauthorization" includes: 1. Patient continues to receive statin at maximally tolerated dose (unless patient has documented inability to take statins), 2. Submission of medical records documenting LDL-C reduction while on Repatha therapy.

# Age Restriction

HeFH,ASCVD: 18 years and older. HoFH: 13 years and older

### **Prescriber Restriction**

## **Coverage Duration**

Initial Authorization will be for 6 months. Reauthorization will be for 1 year.

# **Other Criteria**

A) HoFH: Patient meets one of the following: a) Patient has genetic confirmation of two mutant alleles at LDLR, APOB, PCSK9 or LDLRAP1 gene locus OR b) Patient has an untreated LDL-C level greater than 500 mg/dL (prior to treatment with antihyperlipidemic agents) OR c) Patient has a treated LDL-C level greater than or equal to 300 mg/dL (after treatment with antihyperlipidemic agents but prior to agents such as Repatha, Kynamro, or Juxtapid OR d) Patient has clinical manifestations of HoFH (e.g., cutaneous xanthomas, tendon xanthomas, arcus cornea, tuberous xanthomas or xanthelasma B) HeFH: The patient has a low-density lipoprotein cholesterol (LDL-C level) greater than or equal to 160 mg/dL (after treatment with antihyperlipidemic agents but prior to PCSK9 inhibitor therapy. C) ASCVD: The patient has a low-density lipoprotein cholesterol (LDL-C level) greater than or equal to 160 mg/dL (after treatment with antihyperlipidemic agents but prior to PCSK9 inhibitor therapy. C) ASCVD: The patient has a low-density lipoprotein cholesterol (LDL-C) greater than or equal to 70 mg/dL (after treatment with antihyperlipidemic agents but prior to PCSK9 inhibitor therapy. C) ASCVD: The patient has a low-density lipoprotein cholesterol (LDL-C) greater than or equal to 70 mg/dL (after treatment with antihyperlipidemic agents but prior to PCSK9 inhibitor therapy. ASCVD diagnosis is confirmed by one of the following: acute coronary syndromes, history of myocardial infarction, stable or unstable angina, coronary or other arterial revascularization, stroke, transient ischemic attack, peripheral arterial disease presumed to be of atherosclerotic origin. Examples of high-intensity statin therapy include atorvastatin 40-80 mg, Crestor (rosuvastatin) 20-40 mg. Examples of moderate-intensity statin therapy include atorvastatin 10-20 mg, Crestor (rosuvastatin) 20-40 mg, Examples of moderate-intensity statin therapy include atorvastatin 10-20 mg, Crestor (rosuvastatin) 5-10 mg, simvastatin greater than or equal to 20 mg, pravastatin gre

# Drugs RETROVIR INTRAVENOUS

# **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

**Required Medical Information** Clinical documentation of FDA approved indication for treatment.Clinical documentation of FDA approved indication for treatment.

**Age Restriction** 

**Prescriber Restriction** 

**Coverage Duration** Plan year

Drugs REVLIMID

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

**Exclusion Criteria** Pregnancy (category X)

# **Required Medical Information**

Diagnosis of one of the following: A) Multiple myeloma used in combination with dexamethasone, or B) Diagnosis of transfusion-dependent anemia due to low- or intermediate-1-risk myelodysplastic syndromes associated with a deletion 5q cytogenetic abnormality with or without additional cytogenetic abnormalities, or C) Mantle cell lymphoma and patient's disease has relapsed or progressed after trying at least two prior therapies (Velcade and one of the following: bendamustine, cladribine, fludarabine, rituximab) AND patient is enrolled in the Revlimid REMS Program.

Age Restriction

18 years and older

**Prescriber Restriction** Hematologist/oncologist. Registered in Revlimid REMS.

Coverage Duration

3 Months

#### Drugs REYATAZ ORAL CAPSULE 200 MG, 300 MG, REYATAZ ORAL POWDER IN PACKET

#### **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

**Required Medical Information** Clinical documentation of FDA approved indication for treatment.

#### **Age Restriction**

**Prescriber Restriction** 

**Coverage Duration** Plan year

Drugs RIABNI

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

**Required Medical Information** 

### **Age Restriction**

**Prescriber Restriction** 

**Coverage Duration** 12 months

# **RIBAVIRIN**

# Drugs

ribavirin oral capsule, ribavirin oral tablet 200 mg

# **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

# **Required Medical Information** Clinical documentation of FDA approved indication for treatment.

### **Age Restriction**

**Prescriber Restriction** 

# **Coverage Duration** Per treatment (up to 48 weeks)

#### Drugs RINVOQ

# Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Diagnosis of A) ankylosing spondylitis in adults who have had an inadequate response or intolerance to one or more tumor necrosis factor inhibitors (TNFis), B) Atopic dermatitis for treatment of refractory, moderate to severe atopic dermatitis in patients at least 12 years of age, whose disease is not adequately controlled with other systemic drug products (including biologics) or when those therapies are not advisable, C) Psoriatic arthritis for treatment of active disease in adults who have had an inadequate response or intolerance to one or more TNFis, D) Rheumatoid arthritis for treatment of moderately to severely active disease in adults who have had an inadequate response or intolerance to one or more TNFis OR E) Ulcerative colitis for treatment of moderately to severely active disease in adults who have had an inadequate response or intolerance to one or more TNFis OR E) Ulcerative colitis for treatment of moderately to severely active disease in adults who have had an inadequate response or intolerance to one or more TNFis OR E) Ulcerative colitis for treatment of moderately to severely active disease in adults who have had an inadequate response or intolerance to one or more TNFis

**Age Restriction** 

**Prescriber Restriction** 

**Coverage Duration** 12 months

# Drugs RISPERDAL CONSTA

### **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

### **Required Medical Information**

Clinical documentation of FDA approved indication for treatment. Documented trial of any of the two following drugs: aripiprazole, clozapine, olanzapine, risperidone, quetiapine, ziprasidone.

# Age Restriction

**Prescriber Restriction** 

#### Coverage Duration Plan year

**Other Criteria** Must be unable or unwilling to tolerate oral medications.

#### Drugs ROZLYTREK ORAL CAPSULE 100 MG, 200 MG

# **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

### **Required Medical Information**

Documentation of: A)metastatic non-small cell lung cancer (NSCLC) whose tumors are ROS1-positive OR B) solid tumors that have a neurotrophic tyrosine receptor kinase (NTRK) gene fusion without a known acquired resistance mutation AND are metastatic or where surgical resection is likely to result in severe morbidity AND have either progressed following treatment or have no satisfactory alternative therapy.

Age Restriction 12 years of age or older

Prescriber Restriction Oncology

**Coverage Duration** 3 months

Drugs RUBRACA

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Documented diagnosis of: 1) recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer that had a complete or partial response to platinum-based chemotherapy OR 2) Deleterious BRCA mutation-associated epithelial ovarian, fallopian tube, or primary peritoneal cancer that has been treated with two or more chemotherapies. OR 3)Metastatic Castration-Resistant Prostate Cancer (mCRPC) with BRCA Mutations AND patient has been treated with androgen receptor-directed therapy and a taxane-based chemotherapy.

Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Through end of benefit year

Drugs RYDAPT

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Documentation of: 1) Newly diagnosed acute myeloid leukemia (AML) who are FLT3 mutation-positive, as detected by a FDA approved test and will be used in combination with standard cytarabine and daunorubicin induction and cytarabine consolidation OR 2) Aggressive systemic mastocytosis (ASM), systemic mastocytosis with associated hematological neoplasm (SM-AHN), or mast cell leukemia (MCL)

# **Age Restriction**

**Prescriber Restriction** 

# Coverage Duration

Through end of benefit year

Drugs SABRIL ORAL TABLET, vigabatrin oral powder in packet

# **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

### **Required Medical Information**

Documentation of diagnosis of refractory complex partial seizures (CPS) or infantile spasms (IS). Previously tried and failed two medications for the diagnosis of refractory complex partial seizures including carbamazepine, ethotoin, felbamate, fosphenytoin, gabapentin, lacosamide, lamotrigine, levetiracetam, oxcarbazepine, phenytoin, pregabalin, primidone, tiagabine, topiramate, valproic acid, divalproex sodium, zonisamide.

# Age Restriction

10 years and older for CPS diagnosis. Children aged 1 month to 2 years old for IS.

### Prescriber Restriction

Neurologist registered with the Sabril REMS program

**Coverage Duration** Plan year

Drugs SAMSCA

#### **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

Samsca not approved as an intervention to raise serum sodium urgently to prevent or to treat serious neurological symptoms. Samsca cannot be initiated or re-initiated outside of a hospital setting.

#### **Required Medical Information**

Serum sodium levels. Initial therapy for hyponatremia (hypervolemic and euvolemic): 1. Diagnosis of significant hyponatremia (euvolemic or hypervolemic), AND 2. Treatment has been initiated or re-initiated in a hospital setting prior to discharge. Reauthorization for hypervolemic and euvolemic hyponatremia: 1. Documentation of clinical benefit, AND 2. Treatment hasbeen initiated or re-initiated or re-initiated in a hospital setting prior to discharge.

Age Restriction

**Prescriber Restriction** 

**Coverage Duration** 1 month

**Other Criteria** Documentation of trial and failure of fluid restriction required.

#### Drugs SANDOSTATIN LAR DEPOT INTRAMUSCULAR SUSPENSION, EXTENDED REL RECON

# **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

# **Required Medical Information**

Diagnosis of one of the following: A) Acromegaly and patient had an inadequate response or cannot be treated with surgical resection, pituitary irradiation, and/or bromocriptine mesylate at maximally tolerated doses, or B) Metastatic carcinoid tumor requiring symptomatic treatment of severe diarrhea and flushing episodes, or C) Vasoactive intestinal peptide tumor requiring treatment of profuse watery diarrhea. Acromegaly: Documentation of inadequate response to surgery and/or radiotherapy, or documentation that patient is not a candidate for surgery and/or radiotherapy. Reauthorization will require statement indicating growth hormone (GH) levels are stabilized at less than 5.0ng/mL and IGF-1 levels are normalized (male less than 1.9U/mL or female less than 2.2 U/mL) as matched by age and gender, or the patient has a documented clinical response defined by a reduction of tumor mass, a reduction in the signs and symptoms of acromegaly, or an improvement in significant comorbidities.

# Age Restriction

**Prescriber Restriction** 

Coverage Duration Plan year

### **Other Criteria**

For renewal of acromegaly, IGF-1 level has normalized or improved. For renewal of metastatic carcinoid tumor, patient has improvement in diarrhea and flushing episodes. For renewal of vasoactive intestinal peptide tumor, improvement in diarrhea episodes.

#### Drugs

asenapine maleate, SAPHRIS SUBLINGUAL TABLET 10 MG, 2.5 MG, 5 MG, SECUADO

#### **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

# **Required Medical Information**

Documentation of diagnosis and treatment history

# Age Restriction 18 years and older

### **Prescriber Restriction**

**Coverage Duration** Plan year

#### **Other Criteria**

Member needs to have documented trial of any of the two following drugs: aripiprazole, clozapine, olanzapine, risperidone, quetiapine, ziprasidone

# Drugs SELZENTRY ORAL TABLET 150 MG, 25 MG, 300 MG, 75 MG

**Covered Uses** 

All FDA-approved indications not otherwise excluded by Health Plan.

**Exclusion Criteria** 

#### **Required Medical Information** Clinical documentation of FDA approved indication for treatment.

Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Plan year

**Other Criteria** Documented lab assay verifying HIV is CCR5-tropic positive strain.

# **SEROQUEL XR (Quetiapine)**

#### Drugs

quetiapine oral tablet extended release 24 hr, SEROQUEL XR ORAL TABLET EXTENDED RELEASE 24 HR

# **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

# **Required Medical Information**

Documentation of diagnosis and treatment history

# Age Restriction 10 years and older

# **Prescriber Restriction**

**Coverage Duration** Plan year

# **Other Criteria**

Documentation of reason why quetiapine IR cannot be used.

#### Drugs SIGNIFOR LAR

# **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D. All FDA-approved indications not otherwise excluded from Health Plan.

# **Exclusion Criteria**

# **Required Medical Information**

Documented diagnosis of acromegaly who have had an inadequate response to surgery and/or for whom surgery is not an option

Age Restriction 18 years of age and older

### **Prescriber Restriction**

#### **Coverage Duration** Plan year

#### **Other Criteria**

For renewal, patient's growth hormone level or insulin-like growth factor 1 (IGF-1) level for age and gender has normalized/improved.

# SILDENAFIL (REVATIO)

#### Drugs

sildenafil (pulm.hypertension) oral tablet

### **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

Should not be used in combination with organic nitrates. This product is only indicated for Pulmonary Hypertension and is not to be used for Erectile Dysfunction.

#### **Required Medical Information**

Statement of FDA approved diagnosis of pulmonary arterial hypertension

Age Restriction 18 years and older

**Prescriber Restriction** Cardiologist or Pulmonologist

**Coverage Duration** Plan year

#### Skyrizi

#### Drugs SKYRIZI SUBCUTANEOUS SYRINGE 75 MG/0.83 ML, SKYRIZI SUBCUTANEOUS SYRINGE KIT

# **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Diagnosis of Mod-severe chronic plaque psoriasis (affecting more than 5% of body surface area or crucial body areas such as the hands, feet, face, or genitals) and trial of at least one oral systemic treatment (e.g., MTX, cyclosporine, acitretin, sulfasalazine) OR diagnosis of mod-severe psoriatic arthritis and trial of at least one oral systemic treatment (MTX, etc.) or diagnosis of mod-severe Crohn's disease

# Age Restriction

**Prescriber Restriction** 

**Coverage Duration** 12 months

#### Drugs SOFOSBUVIR-VELPATASVIR

# **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

### **Required Medical Information**

Provider must submit medical records documenting the following: 1) medical diagnosis of Chronic Hepatitis C with labs documenting genotype and subtype, AND 2) medical records documenting viral load taken within 6 months of beginning therapy

# Age Restriction

18 years of age and older

#### **Prescriber Restriction**

Prescribed by, or in consultation with, a gastroenterologist, hepatologist, or infectious disease physician.

### **Coverage Duration**

12 weeks or as defined by current AASLD/IDSA guidance

### **Other Criteria**

Criteria and coverage durations will be applied consistent with current AASLD/IDSA guidance.

#### Drugs SOMATULINE DEPOT

# **Covered Uses**

All FDA-approved indications not otherwise exluded by Health Plan.

# **Exclusion Criteria**

### **Required Medical Information**

Diagnosis of A) acromegaly AND Inadequate response to surgery and/or radiation therapy or patient cannot be treated with surgery and/or radiotherapy, or B) unresectable, well or moderately differentiated, locally advanced or metastatic gastroenteropancreatic neuroendocrine tumors (GEP-NETs).

# Age Restriction

18 years of age and older

# Prescriber Restriction

**Coverage Duration** 3 months initial. Continuation 6 months if no progression

#### **Other Criteria**

For renewal, patient's IGF-1 levels has normalized or improved.

# **SOMAVERT (Pegvisomant)**

#### Drugs

Somavert subcutaneous recon soln 10 mg, SOMAVERT SUBCUTANEOUS RECON SOLN 15 MG, 20 MG, 25 MG, 30 MG

#### **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

#### **Required Medical Information**

Clinical documentation of FDA approved indication for treatment.

Age Restriction 18 years and older

#### **Prescriber Restriction**

Coverage Duration Plan year

Drugs SPRYCEL

**Covered Uses** All FDA-approved indications not otherwise excluded from Health Plan.

# **Exclusion Criteria**

# **Required Medical Information**

Diagnosis of one of the following: A) Newly diagnosed Philadelphia chromosome-positive (Ph+) chronic myeloid leukemia (CML) in chronic phase, or B) Chronic, accelerated, or myeloid or lymphoid blast phase Ph+ CML AND failure, resistance, or intolerance to imatinib, or C) Philadelphia chromosome-positive (Ph+) acute lymphoblastic leukemia (ALL) with resistance or intolerance to imatinib.

Age Restriction 18 years or older

**Prescriber Restriction** Prescriber must be an oncologist.

**Coverage Duration** Plan year

# Stavudine (Zerit)

Drugs

stavudine oral capsule

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

**Required Medical Information** Clinical documentation of FDA approved indication for treatment.

Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Plan year

#### Drugs STIVARGA

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Diagnosis of one of the following: A) Metastatic colorectal cancer AND patient has previously treated with fluoropyrimidine-, oxaliplatin-, and irinotecan-based therapy, an anti-vascular endothelial growth factor (VEGF) therapy, and, if KRAS wild type, an anti-epidermal growth factor receptor (EGFR) therapy, or B) Gastrointestinal stromal tumors that is locally advanced, unresectable or metastatic and patient has tried and had an inadequate response, contraindication or intolerance to Gleevec or Sutent, or C)Hepatocellular carcinoma (HCC) and patient has been previously treated with sorafenib.

Age Restriction 18 years and older

Prescriber Restriction Oncologist

**Coverage Duration** 3 Months

# **Other Criteria**

Hepatic function will be monitored prior to and during treatment and, if patient has elevated liver function tests of hepatocellular necrosis, therapy will be interrupted and then reduced or discontinued.

Drugs STRIBILD

**Covered Uses** All FDA-approved indications not otherwise excluded from Health Plan

**Exclusion Criteria** 

**Required Medical Information** 

**Age Restriction** 

**Prescriber Restriction** 

**Coverage Duration** Plan year

Drugs SUNOSI

**Covered Uses** 

**Exclusion Criteria** 

#### **Required Medical Information**

Clinical documentation of narcolepsy or obstructive sleep apnea AND failed at least TWO alternatives (e.g. methylphenidate, dextroamphetamine, modafinil and armodafinil).

**Age Restriction** 

**Prescriber Restriction** 

**Coverage Duration** Through the plan year

Drugs SUSTIVA

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

**Exclusion Criteria** 

**Required Medical Information** Clinical documentation of FDA approved indication for treatment.

Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Plan year

#### Drugs

sunitinib, SUTENT

#### **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

Combination use with other kinase inhibitors (for example, sorafenib, etc).

# **Required Medical Information**

Diagnosis of one of the following: A) Advanced/metastatic renal cell carcinoma, or B) Gastrointestinal stromal tumors after disease progression on or intolerance to Gleevec, or C) Progressive, well-differentiated pancreatic neuroendocrine tumors in a patient with unresectable locally advanced or metastatic disease.

### **Age Restriction**

Patient must be at least 18 years of age.

# Prescriber Restriction

Must be prescribed by oncologist

# **Coverage Duration**

3 months initial, then renewable in 6 month increments

Drugs SYNERA

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

**Exclusion Criteria** 

**Required Medical Information** Clinical documentation of FDA approved indication for treatment.

Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Plan year

Drugs TABLOID

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

**Required Medical Information** Clinical documentation of FDA approved indication for treatment.

# **Age Restriction**

**Prescriber Restriction** 

**Coverage Duration** Through chemotherapy remission inducation and consolidation treatment

Drugs TAFINLAR

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

### **Required Medical Information**

Diagnosis of unresectable or metastatic melanoma along with BRAF V600E or BRAF V600K mutation status as detected by a US Food and Drug Administration-approved test.

Age Restriction 18 years or older

Prescriber Restriction Oncologist

**Coverage Duration** 3 Months

#### Drugs TAGRISSO ORAL TABLET 40 MG, 80 MG

# **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Documented diagnosis of: 1) metastatic non-small cell lung cancer (NSCLC) with tumors that have epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 L858R mutations, as detected by an FDA-approved test OR 2) metastatic EGFR T790M mutation-positive NSCLC, as detected by an FDA-approved test, with disease progression on or after EGFR tyrosine kinase inhibitor (TKI) therapy.

# Age Restriction

**Prescriber Restriction** 

# Coverage Duration

Through end of benefit year.

#### Takhzyro

#### Drugs TAKHZYRO SUBCUTANEOUS SOLUTION

### **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Documentation that medication is being used to prevent attacks of hereditary angioedema (HAE).

Age Restriction 12 years of age or older

**Prescriber Restriction** 

**Coverage Duration** 3 Months

#### Drugs TALZENNA ORAL CAPSULE 0.25 MG, 1 MG

# **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Documentation of deleterious or suspected deleterious germline breast cancer susceptibility gene (BRCA)-mutated (gBRCAm) human epidermal growth factor receptor 2 (HER2)-negative locally advanced or metastatic breast cancer.

### **Age Restriction**

Prescriber Restriction Oncology

**Coverage Duration** 3 months

#### Drugs ERLOTINIB, TARCEVA

# **Covered Uses**

All FDA approved indications not otherwise excluded by Health Plan. First line for Non-Small Cell Lung Cancer (NSCLC).

# **Exclusion Criteria**

# **Required Medical Information**

Diagnosis of one of the following: A) Locally advanced, unresectable, or metastatic pancreatic cancer and Tarceva will be used in combination with gemcitabine, or B) First-line treatment of metastatic non-small cell lung cancer in which tumors have epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 (L858R) substitution mutations as detected by an Food and Drug Administration (FDA)-approved test, or C) Maintenance treatment of locally advanced or metastatic non-small cell lung cancer when disease has not progressed after 4 cycles of platinum-based first-line chemotherapy, or D) Treatment of locally advanced or metastatic non-small cell lung cancer after failure of at least 1 prior chemotherapy regimen.

# Age Restriction

18 years or older

**Prescriber Restriction** Prescriber must be an oncologist.

Coverage Duration Plan year

# **TARGRETIN** (Bexarotene)

#### Drugs

bexarotene oral, TARGRETIN TOPICAL

# **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

**Required Medical Information** Documentation of diagnosis and treatment history.

# Age Restriction 18 years and older.

# **Prescriber Restriction**

**Coverage Duration** Plan year

#### Drugs TASIGNA ORAL CAPSULE 150 MG, 200 MG

# **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

### **Exclusion Criteria**

Long QT syndrome. Uncorrected hypokalemia. Uncorrected hypomagnesemia. Concomitant use with a drug known to prolong the QT interval or strong cytochrome P450 3A4 inhibitors.

#### **Required Medical Information**

Diagnosis of one of the following: A) Newly diagnosed adult patients with Philadephia chromosome positive chronic myeloid leukemia (Ph+ CML) in chronic phase, or B) Ph+ chronic or accelerated phase chronic myeloid leukemia (CML) in adult patients resistant to or intolerant to prior therapy that included imatinib.

Age Restriction 18 years and older

**Prescriber Restriction** Must be prescribed by Oncologist

**Coverage Duration** Initial: 3 months, Renewal: 6 months with documentation of continued benefit

#### Drugs TAVALISSE

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Documented Platelet count less than 30x10^9/L and member had an insufficient response to previous treatment (corticosteroids, immunoglobulins, splenectomy, and/or a thrombopoietin receptor agonists)

# Age Restriction 18 years or older

#### Prescriber Restriction Hematologist

**Coverage Duration** Initial Authorization will be for 3 months. Reauthorization will be for 1 year.

Drugs TAZVERIK

**Covered Uses** 

**Exclusion Criteria** 

# **Required Medical Information**

Documentation of metastatic or locally advanced epithelioid sarcoma that is not eligible for complete resection.

Age Restriction

Prescriber Restriction Oncology

**Coverage Duration** 3 months

### Drugs

dimethyl fumarate oral capsule, delayed release (DR/EC) 120 mg, 120 mg (14)- 240 mg (46), 240 mg

# Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

### **Exclusion Criteria**

Treatment of primary progressive MS is not covered. Combination therapy with a beta interferon product, Gilenya, Aubagio, Tecfidera, Tysabri or Copaxone is not covered.

#### **Required Medical Information**

Diagnosis of relapsing forms of multiple sclerosis (relapsing-remitting MS or progressive-relapsing MS, or secondaryprogresive MS) OR patient has experienced a first clinical episode and has MRI features consistent with multiple sclerosis.

**Age Restriction** 

**Prescriber Restriction** 

**Coverage Duration** Plan year

Drugs

temozolomide

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

#### **Required Medical Information** Clinical documentation of FDA approved indication for treatment.

Age Restriction 18 years and older

**Prescriber Restriction** 

**Coverage Duration** 3 Months

Drugs TEPMETKO

**Covered Uses** 

**Exclusion Criteria** 

**Required Medical Information** Diagnosis of Non-small cell lung cancer, metastatic (with mesenchymal-epithelial transition [MET] exon 14 skipping mutation).

Age Restriction 18 years of age and older

**Prescriber Restriction** Oncology

Coverage Duration Through end of benefit year

Drugs THALOMID

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

Pregnancy (category X)

# **Required Medical Information**

Diagnosis of one of the following: A) Multiple myeloma that is newly diagnosed and is receiving concurrent dexamethasone, or B) Acute treatment of cutaneous manifestations of moderate to severe erythema nodosum leprosum (ENL) AND the medication will not be used as monotherapy if the member has moderate to severe neuritis, or C) Maintenance therapy for prevention and suppression of cutaneous manifestations of ENL recurrence.

Age Restriction

12 years of age and older

### **Prescriber Restriction**

Candidates must follow Thalomid REMS program requirements. Provider and pharmacy must be registered with this program.

**Coverage Duration** 3 months

# Drugs THYMOGLOBULIN

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

# **Required Medical Information** Clinical documentation of FDA approved indication for treatment.

Age Restriction 18 years and older

# **Prescriber Restriction**

**Coverage Duration** Up to 14 days.

Drugs TIBSOVO

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Documentation of relapsed or refractory acute myeloid leukemia (AML) with a susceptible isocitrate dehydrogenase-1 (IDH1) mutation as detected by an FDA-approved test.

# Age Restriction

Prescriber Restriction Oncology

**Coverage Duration** 3 months

# Drugs TIGAN INTRAMUSCULAR

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

# **Required Medical Information** Clinical documentation of FDA approved indication for treatment.

Age Restriction Use not recommended in children

# **Prescriber Restriction**

Coverage Duration Per treatment

Drugs TIVICAY

**Covered Uses** All FDA-approved indications not otherwise excluded from Health Plan

**Exclusion Criteria** 

**Required Medical Information** 

**Age Restriction** 

**Prescriber Restriction** 

**Coverage Duration** Plan year

# **Tobramycin Inhalant Solution (TOBI)**

Drugs

tobramycin in 0.225 % NaCl

**Covered Uses** All FDA-approved indications not otherwise excluded from Health Plan.

**Exclusion Criteria** 

**Required Medical Information** Documented diagnosis of cystic fibrosis with Pseudomonas

Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Plan yer

Drugs TOPOSAR

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

**Exclusion Criteria** 

**Required Medical Information** Must be prescribed for an FDA approved indication

# Age Restriction

Prescriber Restriction Oncologist

**Coverage Duration** 12 months

#### Drugs

bosentan, TRACLEER

#### **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

Receiving concomitant cyclosporine A or glyburide therapy. Aminotransferase elevations are accompanied by signs or symptoms of liver dysfunction or injury or increases in bilirubin at least 2 times the upper limit of normal. For female patients, pregnancy must be excluded prior to the start of therapy and will be prevented thereafter with reliable contraception.

#### **Required Medical Information**

Diagnosis of pulmonary arterial hypertension (PAH) WHO Group I AND New York Heart Association (NYHA) Functional Class II-IV

**Age Restriction** Greater than 12 years of age

### **Prescriber Restriction**

Available only to those enrolled in the Tracleer REMS Program. Prescription is written by or in consultation with a pulmonologist or cardiologist

# **Coverage Duration**

3 Months

#### **Other Criteria**

Liver aminotransferases will be measured prior to initiation of treatment and then monthly.

# Tranexamic acid (CYKLOKAPRON)

#### Drugs

tranexamic acid intravenous

### **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

**Required Medical Information** Documentation of hemophilia diagnosis as appropriate

# Age Restriction

**Prescriber Restriction** 

**Coverage Duration** 8 days

Drugs

ALTRENO, tretinoin (antineoplastic)

# **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

# **Required Medical Information**

Patients diagnosed with acne vulgaris after trying and failing at least 1 preferred alternatives (such as generic acne products - erythromycin/benzoyl peroxide, clindamycin, etc) or other non-cosmetic diagnosis.

# Age Restriction

**Prescriber Restriction** 

**Coverage Duration** 3 months

#### Trikafta

#### Drugs TRIKAFTA ORAL TABLETS, SEQUENTIAL 100-50-75 MG(D) /150 MG (N)

# **Covered Uses**

**Exclusion Criteria** Use in combination with other CFTR modulator (Orkambi, Kalydeco, or Symdeko)

# **Required Medical Information**

Documentation of Cystic Fibrosis AND confirmation of presence of at least one F508del mutation in CFTR gene through genetic testing.

Age Restriction 12 years of age or older

# **Prescriber Restriction**

Prescribed by pulmonologist or a physician who specializes in the treatment of Cystic fibrosis

# **Coverage Duration**

Through end of benefit year

Drugs TRINTELLIX

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Diagnosis of major depressive disorder and treatment failure of at least two other formulary medications used in the treatment of major depressive disorder

# Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Through end of benefit year.

Drugs TRIUMEQ

**Covered Uses** All FDA-approved indications not otherwise excluded from Health Plan

**Exclusion Criteria** 

**Required Medical Information** 

Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Plan year

Drugs TRUVADA

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

**Exclusion Criteria** 

**Required Medical Information** Clinical documentation of FDA approved indication for treatment.

Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Plan year

Drugs TURALIO

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Documentation of symptomatic tenosynovial giant cell tumor (TGCT) associated with severe morbidity or functional limitations and not amenable to improvement with surgery.

# Age Restriction

Prescriber Restriction Oncologist

**Coverage Duration** 3 months

#### Drugs

lapatinib, TYKERB

#### **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

### **Required Medical Information**

Diagnosis of breast cancer with tumors that overexpress human epidermal growth factor receptor 2 (HER2) AND 1. the medication will be used in combination with Xeloda in a patient with advanced or metastatic disease and the patient has received prior therapy including an anthracycline, a taxane, and trastuzumab, or 2) The medication will be used in combination with Femara for the treatment of a postmenopausal woman with hormone receptor-positive metastatic disease for whom hormonal therapy is indicated.

### **Age Restriction**

18 years or older

### **Prescriber Restriction**

Must be prescribed by Oncologist and Oncologist must monitor treatment

#### **Coverage Duration**

Initial: 3 months, Renewal: 6 months with documentation of continued benefit

Drugs TYMLOS

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Diagnosis of one of the following: A) Osteoporosis in a postmenopausal female AND one or more of the following: 1) History of osteoporotic fracture, or 2) Documented trial and failure of bisphosphonate or 3) Documented contraindication or intolerance to bisphosphonate therapy. Patient has not received more than 2 years of therapy with Tymlos.

# **Age Restriction**

# **Prescriber Restriction**

# **Coverage Duration**

Initial: 1 year. Renewal: 1 year not to exceed 2 years of total

Drugs UBRELVY

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Documentation that the medication will be used for the acute treatment of migraine with or without aura in adults AND the member has tried and failed two triptans, unless contraindicated.

# Age Restriction

**Prescriber Restriction** 

#### **Coverage Duration** Through end of benefit year

**Other Criteria** 

Reauthorization requires documentation of medication efficacy.

# Drugs VALCYTE ORAL RECON SOLN, valganciclovir oral tablet

# **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

# **Required Medical Information**

Clinical documentation of FDA approved indication for treatment.

Age Restriction 4 months and older.

# **Prescriber Restriction**

**Coverage Duration** Plan year

#### Drugs

vancomycin oral capsule

#### **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

### **Exclusion Criteria**

### **Required Medical Information**

Diagnosis of A) Clostridium difficile-associated diarrhea, AND Stool culture report within the previous 30 days indicating positive C. difficile toxin, AND documented trial and failure or contraindication to preferred agent, metronidazole, or B) Staphylococcus aureus (including methicillin-resistant strains)enterocolitis

### Age Restriction

### **Prescriber Restriction**

### **Coverage Duration**

14 days, Patients with multiple relapses: 6 weeks

#### Drugs

calcitriol topical

#### **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

### **Exclusion Criteria**

Any pertinent clinical situation as defined by the product label that could affect patient safety and/or therapeutic efficacy (i.e. contraindications, warnings, precautions, adverse effects, renal or hepatic function, drug interactions, lab values, required prior or concomitant therapy, inappropriate dosing and/or duration, etc).

### **Required Medical Information**

Clinical documentation of FDA approved indication for treatment.

**Age Restriction** 

**Prescriber Restriction** 

**Coverage Duration** Plan year

#### **Other Criteria**

Must have tried and failed at least 2 topical steroids (at least one mid potency and at least one high potency)

#### Drugs VENCLEXTA, VENCLEXTA STARTING PACK

# **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

### **Required Medical Information**

Documented diagnosis of A) chronic lymphocytic leukemia (CLL) or small lymphocyctic lymphoma (SLL) or B) newlydiagnosed acute myeloid leukemia (AML) in adults who are age 75 years or older, or who have comorbidities that preclude use of intensive induction chemotherapy and will be used in combination with azacitidine, decitabine, or lowdose cytarabine.

# Age Restriction

**Prescriber Restriction** 

**Coverage Duration** 3 months

Drugs VENTAVIS

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

### **Required Medical Information**

Diagnosis of Pulmonary Arterial Hypertension (PAH) AND has WHO Group I PAH AND Patient has New York Heart Association (NYHA) Functional Class III or IV. Trial and failure of Revatio or Adcirca.

#### Age Restriction 18 years or older

To years of older

# **Prescriber Restriction**

Must be prescribed by a cardiologist or pulmonologist

# **Coverage Duration** 3 months.

Drugs VERQUVO

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

#### **Exclusion Criteria**

#### **Required Medical Information**

Heart failure, chronic, To reduce the risk of cardiovascular death and heart failure hospitalization in symptomatic patients with reduced ejection fraction

# Age Restriction

18 years or older

#### Prescriber Restriction Cardiologist

**Coverage Duration** 

1 year

#### **Other Criteria**

All the below must be met. 1) Diagnosis of chronic heart failure of NYHA class II-IV 2)Must have LVEF less than 45% 3)Concomitant or prior treatment with Ace inhibitor/Angiotensin receptor blocker/Entresto and a beta blocker 4) Dose does not exceed 10mg/day 5)Must have had a previous hospitalization due to heart failure within the past 6 months, or an outpatient intravenous diuretic treatment for heart failure within the past 3 months.

Drugs VERZENIO

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

#### **Exclusion Criteria**

#### **Required Medical Information**

Documented diagnosis of one of the following: A)hormone receptor -positive, human epidermal growth factor receptor 2 - negative advanced or metastatic breast cancer and must ne used in combination with fulvestrant unless there is disease progression following endocrine therapy and prior chemotherapy in the metastatic setting.

#### Age Restriction

Prescriber Restriction Oncologist

**Coverage Duration** 3 Months

#### Drugs VIBERZI

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Clinically diagnosed with irritable bowel syndrome with diarrhea supported by documentation from the patient's medical records AND Other GI medical conditions that could explain the symptoms have been ruled out AND Failed conventional non-pharmacological therapies including (Dietary changes, stress reduction, or behavioral changes)AND Failed conventional pharmacological therapies including: Antidiarrheals, Antidepressants, and Antispasmodics AND Must have tried and failed rifaximin

# Age Restriction

#### **Prescriber Restriction**

# **Coverage Duration**

Initial Authorization will be for 3 months. Reauthorization will be for 1 year.

#### Drugs VIIBRYD ORAL TABLET, VIIBRYD ORAL TABLETS,DOSE PACK 10 MG (7)- 20 MG (23)

#### **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D.

#### **Exclusion Criteria**

# **Required Medical Information**

Documented trial of at least two generic antidepressants (e.g. bupropion, citalopram, escitalopram, fluoxetine, fluvoxamine, paroxetine, paroxetine CR, sertraline, duloxetine, venlafaxine).

#### Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Through end of benefit year

Drugs VILTEPSO

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Diagnosis of Duchenne muscular dystrophy by, or in consultation with, a neurologist with expertise in the diagnosis of DMD AND Submission of medical records (e.g., chart notes, laboratory values) confirming the mutation of the DMD gene amenable to exon 53 skipping AND One of the following 1) patient has a 6-Minute OR Walk Time (6MWT) greater than 300 meters while walking independently (e.g., without side-by-side assist, cane, walker, wheelchair, etc.) prior to beginning Viltepso therapy OR 2) the patient is ambulatory without needing an assistive device AND has achieved a score of greater than 17 on the North Star Ambulatory Assessment (NSAA) AND dosing for DMD is in accordance with the United States Food and Drug Administration approved labeling

# Age Restriction

# **Prescriber Restriction**

Prescribed by, or in consultation with, a neurologist with expertise in the treatment of DMD

**Coverage Duration** 6 months

**Other Criteria** Reauthorization: Documentation of an increase in the physical function from baseline

# Drugs VIMPAT ORAL SOLUTION, VIMPAT ORAL TABLET, VIMPAT ORAL TABLETS, DOSE PACK

#### **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

#### **Required Medical Information**

Age Restriction 17 years of age or older

# **Prescriber Restriction**

**Coverage Duration** Through the benefit year

Other Criteria Max dose 400mg/day

# Drugs VIRACEPT ORAL TABLET

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

# Exclusion Criteria

**Required Medical Information** Clinical documentation of FDA approved indication for treatment.

# Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Plan year

#### **VIRAMUNE** (Nevirapine)

**Drugs** nevirapine

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

**Required Medical Information** Clinical documentation of FDA approved indication for treatment.

#### Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Plan year

Drugs VIREAD

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

**Exclusion Criteria** 

**Required Medical Information** Clinical documentation of FDA approved indication for treatment.

Age Restriction

**Prescriber Restriction** 

Coverage Duration Plan year

### VITEKTA (Elvitegravir)

Drugs tenofovir disoproxil fumarate

**Covered Uses** All FDA-approved indications not otherwise excluded from Health Plan

#### **Exclusion Criteria**

**Required Medical Information** 

# Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Plan year

#### Drugs VITRAKVI ORAL CAPSULE

# Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

#### **Exclusion Criteria**

# **Required Medical Information**

Clinical documentation of unresectable or metastatic solid tumors that have a neurotrophic receptor tyrosine kinase (NTRK) gene fusion and have no satisfactory alternative treatments or that have progressed following treatment.

#### **Age Restriction**

Prescriber Restriction Oncology

**Coverage Duration** 3 months

Drugs VIZIMPRO

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

#### **Required Medical Information**

Documentation that medication will be used for the first-line treatment of patients with metastatic non-small cell lung cancer (NSCLC) with epidermal growth factor receptor (EGFR) exon 19 deletion or exon 21 L858R substitution mutations as detected by an FDA-approved test.

#### Age Restriction

Prescriber Restriction Oncology

**Coverage Duration** 3 months

#### Drugs

voriconazole oral

#### **Covered Uses**

All medically-accepted indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

#### **Required Medical Information**

Documentation of invasive aspergillosis, bronchopulmonary aspergillosis, candidemia and disseminated candidiasis in skin, abdomen, kidney, bladder wall and wounds, esophageal candidiasis, and serious Candida infections, infections caused by the emerging pathogens Scedosporium sp. and Fusarium sp., or rare and refractory fungal infections should be provided. Preferred alternative for Candida: oral fluconazole

Age Restriction 12 years or older

**Prescriber Restriction** 

**Coverage Duration** 6 months

Drugs VOTRIENT

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

#### **Required Medical Information**

Diagnosis of one of the following: A) Advanced/metastatic renal cell carcinoma, or B) Advanced soft tissue sarcoma and patient received at least one prior chemotherapy (e.g., doxorubicin, dacarbazine, ifosfamide, epirubicin, gemicitabine, docetaxel, or vinorelbine).

Age Restriction 18 years of age and older

Prescriber Restriction Oncologist

**Coverage Duration** Initial: 3 months, Renewal: 6 months with documentation of continued benefit

#### Drugs VUMERITY

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

Concurrent use with other MS disease modifying agents (including Aubagio, Avonex, Betaseron, Copaxone/Glatiramer/Glatopa, Extavia, Gilenya, Lemtrada, Mavenclad, Mayzent, Ocrevus, Plegridy, Rebif, Tecfidera and Tysabri) OR Member with moderate or severe renal impairment (creatinine clearance less than 60 mL/min.

# **Required Medical Information**

Diagnosis of relapsing forms of multiple sclerosis and must have tried and failed one generic alternative agent such as dimethyl fumarate or glatiramer acetate

Age Restriction

Prescriber Restriction Neurologist

Coverage Duration Benefit Year

#### Drugs VYVANSE ORAL CAPSULE

#### **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D.

#### **Exclusion Criteria**

#### **Required Medical Information**

Diagnosis of one of the following: A) ADHD and tried and failed two alternative medications FDA approved for the treatment of ADHD, or B) Moderate to severe Binge Eating Disorder AND the patient is receiving psychological counseling AND the patient must have tried and failed at least two antidepressant medications.

#### Age Restriction

ADHD: Must be older than 6 years of age, BED: Must be 18 years of age or older

#### **Prescriber Restriction**

BED: The medication must be prescribed by a psychiatrist or a psychiatric specialist.

Coverage Duration

12 months

Drugs VYXEOS

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

#### **Exclusion Criteria**

Documented diagnosis of newly-diagnosed therapy-related acute myeloid leukemia (t-AML) or AML with myelodysplasiarelated changes (AML-MRC).

### **Required Medical Information**

#### **Age Restriction**

Prescriber Restriction oncologist or hematologist

**Coverage Duration** 3 Months

Drugs WAKIX

**Covered Uses** 

**Exclusion Criteria** 

#### **Required Medical Information**

Diagnosis of one of Excessive daytime sleepiness in patients with narcolepsy AND 1. Submission of sleep study with narcolepsy diagnosis, AND 2. currently NOT taking any sedative hypnotics or other CNS depressants AND 3. has experienced inadequate response or intolerable side effects to two preferred products (modafinil, armodofinil, methylphenidate, or dextroamphetamine).

#### **Age Restriction**

#### **Prescriber Restriction**

#### **Coverage Duration**

Initial: 3 months, Renewal: 6 months with documentation of continued benefit

#### **Other Criteria**

For renewal, the patient had a positive response to the medication (increased sleep quality for patients with narcolepsy).

Drugs XALKORI

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

#### **Required Medical Information**

Stated diagnosis of late-stage (locally advanced or metastatic), non-small cell lung cancers (NSCLC) with expression of the abnormal anaplastic lymphoma kinase (ALK) gene as detected by an FDA approved test.

Age Restriction 18 years and older

Prescriber Restriction Oncologist

**Coverage Duration** 3 months

#### Drugs

# XCÕPRI, XCOPRI MAINTENANCE PACK ORAL TABLET 250MG/DAY(150 MG X1-100MG X1), 350 MG/DAY (200 MG X1-150MG X1), XCOPRI TITRATION PACK

#### **Covered Uses**

**Exclusion Criteria** 

#### **Required Medical Information**

Diagnosed with partial-onset seizures and treatment failure of at least two other formulary medications used in the treatment of partial onset seizures from the following list: Briviact, clobazam, felbamate, lamotrigine, levetiracetam, Fycompa, rufinamide, topiramate, valproate, zonisamide, carbamazepine, Aptiom, gabapentin, Vimpat, oxcarbazepine, phenobarbital, phenytoin, pregabalin, primidone, tiagabine, vigabatrin.

#### **Age Restriction**

**Prescriber Restriction** 

#### Coverage Duration

Through end of benefit year

#### Drugs XELJANZ ORAL SOLUTION, XELJANZ ORAL TABLET, XELJANZ XR

# **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Diagnosis of one of the following, A) Ankylosing spondylitis in adults with active disease who have had an inadequate response or intolerance to one or more tumor necrosis factor inhibitors (TNFis) B) Polyarticular juvenile idiopathic arthritis (JIA), in patients at least 2 years of age with active disease who have had an inadequate response or intolerance to one or more TNFis C) Psoriatic Arthritis in adults with active disease who have had an inadequate response or intolerance to one or more TNFis D) Rheumatoid arthritis in adults with moderately to severely active disease who have had an inadequate response or intolerance to one or more TNFis OR E) Ulcerative colitis in adults with moderately to severely active disease who have had an inadequate response or intolerance to one or more TNFis OR E) Ulcerative colitis in adults with moderately to severely active disease who have had an inadequate response or who are intolerant to one or more TNFis

#### Age Restriction

**Prescriber Restriction** 

**Coverage Duration** 12 months

#### Drugs XIFAXAN ORAL TABLET 200 MG, 550 MG

# **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

Allergy to rifamycin agents

#### **Required Medical Information**

Diagnosis of traveler's diarrhea and patient does not have fever or blood in the stool OR Diagnosis of hepatic encephalopathy and tried and failed lactulose therapy OR Diagnosis of irritable bowel syndrome with diarrhea (IBS-D) and failed at least TWO alternatives from TWO different classes such as anitdiarrheals (e.g. loperamide, diphenoxylateatropine), antispasmodics (e.g. dicyclomine), bile acid sequestrants (e.g. cholestryramine, colestipol).

#### **Age Restriction**

Traveler's diarrhea: 12 years of age or older, Hepatic encephalopathy and IBS-D: 18 years of age or older

#### **Prescriber Restriction**

**Coverage Duration** Traveler's diarrhea: 3 days, Hepatic encephalopathy: 6 months, IBS-D: 6 weeks

Drugs XOLAIR

#### **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

#### **Required Medical Information**

Diagnosis of A) moderate to severe persistent allergic asthma AND Evidence of specific allergic sensitivity confirmed by positive skin test (i.e. prick/puncture test) or blood test (i.e. radioallergosorbent test) for a specific IgE or in vitro reactivity to a perennial aeroallergen AND Pretreatment serum IgE levels greater than 30 and less than 700 IU/mL AND Symptoms are not adequately controlled with at least ONE inhaled corticosteroid (ICS) plus long-acting beta2-agonist (LABA) (e.g. Advair, Symbicort) for at least 3 months unless patient is intolerant to such treatment or such treatment is contraindicated, or B) Treatment of chronic idiopathic urticaria in adults and adolescents 12 years and older who remain symptomatic despite H1 antihistamine treatment (i.e. loratidine, cetirizine, levocetirizine, fexofenadine, etc.).

#### **Age Restriction**

Patient must be 12 years of age or older

#### **Prescriber Restriction**

Initial drug order must be by an allergist/immunologist, dermatologist, or a pulmonologist

**Coverage Duration** 6 months

Drugs XOSPATA

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

#### **Exclusion Criteria**

# **Required Medical Information**

Clinical documentation of relapsed or refractory acute myeloid leukemia (AML) with a FMS-like tyrosine kinase 3 (FLT3) mutation as detected by an FDA-approved test.

#### Age Restriction

Prescriber Restriction Oncology

**Coverage Duration** 3 months

#### Drugs XPOVIO ORAL TABLET 60MG TWICE WEEK (120 MG/WEEK), 80MG TWICE WEEK (160 MG/WEEK)

#### **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D.

#### **Exclusion Criteria**

#### **Required Medical Information**

Documentation that the medication will be used in 1) combination with dexamethasone for the treatment of adult patients with relapsed or refractory multiple myeloma (RRMM) who have received at least four prior therapies and whose disease is refractory to at least two proteasome inhibitors, at least two immunomodulatory agents, and an anti-CD38 monoclonal antibody. OR 2)monotherapy for the treatment of adult patients with relapsed or refractory diffuse large B-cell lymphoma

#### Age Restriction

Prescriber Restriction Oncologist

**Coverage Duration** 3 months

#### Drugs XTANDI ORAL CAPSULE

#### Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

#### **Required Medical Information**

Diagnosis of metastatic castration-resistant prostate cancer AND the patient has tried and had an inadequate response, contraindication or intolerance to Zytiga.

#### Age Restriction

Prescriber Restriction Oncologist or urologist

# **Coverage Duration** 3 months

**Other Criteria** Must try and fail Zytiga first.

#### Drugs XYREM

# **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

Contraindications: Patient is being treated with sedative hypnotic agents, other CNS depressants, or using alcohol. Patient has succinic semialdehyde dehydrogenase deficiency. Patient has a history of drug abuse.

# **Required Medical Information**

Diagnosis of one of the following: A) Excessive daytime sleepiness in patients with narcolepsy AND 1. Submission of sleep study with narcolepsy diagnosis, AND 2. currently NOT taking any sedative hypnotics or other CNS depressants AND 3. has experienced inadequate response or intolerable side effects to two preferred products (modafinil, armodofinil, methylphenidate, or dextroamphetamine) AND 4. The requested dose does not exceed the FDA indicated maximum (9gm/night), or B) Cataplexy in patients with narcolepsy AND 1. Submission of sleep study showing narcolepsy diagnosis, AND 2. currently NOT taking any sedative hypnotics or other CNS depressants, AND 3. does not have sleep apnea, AND 4. The dose does not exceed the FDA indicated maximum (9gm/night).

# Age Restriction

18 years of age and older

# Prescriber Restriction

Neurologist

#### **Coverage Duration**

Initial: 3 months, Renewal: 6 months with documentation of continued benefit

#### **Other Criteria**

Patient and physician must adhere to all regulations of the Xyrem REMS Program. For renewal, the patient had a positive response to the medication (increased sleep quality for patients with narcolepsy).

Drugs YONSA

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

#### **Exclusion Criteria**

#### **Required Medical Information**

Documented diagnosis of metastatic castration resistant prostate cancer (CRPC)AND the medication is being used in combination with methylprednisolone AND medication not being used as dual therapy with another androgen receptor inhibitor.

Age Restriction 18 years of age or older

# **Prescriber Restriction**

**Coverage Duration** 3 Months

Drugs ZAVESCA

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

### **Exclusion Criteria**

# **Required Medical Information**

Medical statement of approved diagnosis: mild to moderate type 1 Gaucher disease for whom enzyme replacement therapy is not a therapeutic option (e.g., due to constraints such as allergy, hypersensitivity, or poor venous access).

Age Restriction 18 years or older

**Prescriber Restriction** 

Coverage Duration Plan year

Drugs ZEJULA

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

# **Exclusion Criteria**

# **Required Medical Information**

Documented diagnosis of 1) recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer in a complete or partial response to platinum-based chemotherapy OR 2) advanced epithelial ovarian, fallopian tube, or primary peritoneal cancer who are in a complete or partial response to first-line platinum-based chemotherapy OR 3) advanced ovarian, fallopian tube, or primary peritoneal cancer who have been treated with three or more prior chemotherapy regimens and whose cancer is associated with homologous recombination deficiency (HRD) positive status.

# Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Through end of benefit year

Drugs ZELBORAF

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

#### **Exclusion Criteria** Combination use with ipilimumab

# **Required Medical Information**

Diagnosis of A) unresectable or metastatic melanoma whose tumors express a gene mutation called BRAF V600E detected by an FDA approved test. Vemurafenib is not recommended for use in patients with wild-type BRAF melanoma. OR B) Erdheim-Chester Disease (ECD) with BRAF V600 mutation.

Age Restriction 18 years or older

Prescriber Restriction Oncologist

**Coverage Duration** 3 months

#### Drugs ZEMAIRA

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

# Required Medical Information

Clinical documentation of FDA approved indication for treatment.

Age Restriction 18 years and older.

**Prescriber Restriction** 

**Coverage Duration** 3 months.

#### **ZIAGEN** (Abacavir)

#### Drugs

#### abacavir oral tablet, ZIAGEN ORAL SOLUTION

#### **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

**Required Medical Information** Clinical documentation of FDA approved indication for treatment.

#### **Age Restriction**

**Prescriber Restriction** 

**Coverage Duration** Plan year

**Other Criteria** Documented negative HLA-B\*5701 screening.

#### Zidovudine (RETROVIR)

Drugs zidovudine

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

**Required Medical Information** Clinical documentation of FDA approved indication for treatment.

#### Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Plan year

#### Drugs

zoledronic acid-mannitol-water intravenous piggyback 5 mg/100 mL

#### **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

#### **Exclusion Criteria**

#### **Required Medical Information**

Clinical documentation of FDA approved indication for treatment. For treatment of hypercalcemia of malignancy, must have documentation of corrected total serum calcium greater than or equal to 12 mg/dL. For Paget's disease, must have symptomatic form of disease.

# Age Restriction 18 years and older

# Prescriber Restriction

# Coverage Duration

Per treatment

#### **Other Criteria**

For Paget's disease, must have documented failure, intolerance or contraindication to oral agent: alendronate OR risedronate. For osteoporosis, must have documented failure, intolerance or contraindication to at least 2 oral bisphosphonates.

# Drugs ZOLINZA

**Covered Uses** All FDA-approved indications not otherwise excluded by Health Plan.

# **Exclusion Criteria**

# **Required Medical Information**

Clinical documentation of FDA approved indication for treatment.

Age Restriction 18 years and older.

# **Prescriber Restriction**

**Coverage Duration** Plan year

#### **Other Criteria**

Treatment may be continued as long as there is no evidence of progressive disease or unacceptable toxicity.

#### Drugs

everolimus (immunosuppressive) oral tablet 0.25 mg, 0.5 mg, 0.75 mg, **ZORTRESS ORAL TABLET 0.25 MG, 0.5 MG**, **0.75 MG** 

## **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

## **Exclusion Criteria**

Any pertinent clinical situation as defined by the product label that could affect patient safety and/or therapeutic efficacy (i.e. contraindications, warnings, precautions, adverse effects, renal or hepatic function, drug interactions, lab values, required prior or concomitant therapy, inappropriate dosing and/or duration, etc).

#### **Required Medical Information**

Clinical documentation of FDA approved indication for treatment.

Age Restriction 18 years and older.

**Prescriber Restriction** 

**Coverage Duration** 6 months.

## ZYDELIG

Drugs ZYDELIG

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

**Exclusion Criteria** 

**Required Medical Information** 

Age Restriction

**Prescriber Restriction** 

**Coverage Duration** Through end of benefit year.

Drugs ZYFLO

## **Covered Uses**

All FDA-approved indications not otherwise excluded from Part D.

#### **Exclusion Criteria**

Any pertinent clinical situation as defined by the product label that could affect patient safety and/or therapeutic efficacy (i.e. contraindications, warnings, precautions, adverse effects, renal or hepatic function, drug interactions, lab values, required prior or concomitant therapy, inappropriate dosing and/or duration, etc).

#### **Required Medical Information**

Documentation of diagnosis and treatment history.

Age Restriction 12 years and older.

**Prescriber Restriction** 

**Coverage Duration** Plan year

**Other Criteria** Must have failed montelukast and zafirlukast.

## ZYKADIA

### Drugs ZYKADIA ORAL TABLET

**Covered Uses** All FDA-approved indications not otherwise excluded from Part D.

## **Exclusion Criteria**

## **Required Medical Information**

Documented diagnosis of metastatic non-small cell lung cancer (NSCLC) whose tumors are anaplastic lymphoma kinase (ALK)-positive as detected by an FDA-approved test

## Age Restriction

**Prescriber Restriction** 

**Coverage Duration** 3 months

#### Drugs

abiraterone oral tablet 250 mg, 500 mg, ZYTIGA ORAL TABLET 250 MG, 500 MG

## **Covered Uses**

All FDA-approved indications not otherwise excluded by Health Plan.

## **Exclusion Criteria**

## **Required Medical Information**

Diagnosis of metastatic prostate cancer AND Patient has castration-resistant disease (defined by tumor growth/disease progression, risk in PsA levels, new metastases) AND Zytiga will be used in combination with prednisone.

## Age Restriction

**Prescriber Restriction** 

Coverage Duration 3 months

# Index

abacavir oral tablet	
abacavir-lamivudine	. 82
abiraterone oral tablet 250 mg, 500	
mg	329
acitretin	
ACTIMMUNE	
adefovir	
ADEMPAS	
AFINITOR ORAL TABLET 10 MG,	
2.5 MG, 5 MG, 7.5 MG	7
AIMOVIG AUTOINJECTOR	
AJOVY AUTOINJECTOR	
AJOVY AUTOINJECTOR	
	.43
AKYNZEO (FOSNETUPITANT)	~~
INTRAVENOUS RECON SOLN	
AKYNZEO (NETUPITANT)	
ALECENSA	
ALINIA	
ALIQOPA	
alosetron	
ALTRENO	275
ALUNBRIG ORAL TABLET 180	
MG, 30 MG, 90 MG	.12
ALUNBRIG ORAL	
TABLETS, DOSE PACK	. 12
ambrisentan	
APTIOM	
APTIVUS	
ARCALYST	
armodafinil	
asenapine maleate	
ATRIPLA	
AUBAGIO	
AUSTEDO ORAL TABLET 12 MG,	
6 MG, 9 MG	
AVONEX INTRAMUSCULAR PEN	20
	04
INJECTOR AVONEX INTRAMUSCULAR PEN	. 2 I
	~ 4
INJECTOR KIT AVONEX INTRAMUSCULAR	.21
	~ (
SYRINGE	.21
AVONEX INTRAMUSCULAR	
SYRINGE KIT	. 21
AYVAKIT ORAL TABLET 100	
MG, 200 MG, 300 MG	
BALVERSA	
BANZEL ORAL SUSPENSION	.24
BANZEL ORAL TABLET 200 MG,	
400 MG	
BARACLUDE ORAL SOLUTION	25
BAVENCIO	. 26
BELSOMRA	27
BENLYSTA SUBCUTANEOUS	
AUTO-INJECTOR	28
benznidazole	
BETASERON SUBCUTANEOUS	-
KIT	31
bexarotene oral	
BLENREP	

BOSULIF ORAL TABLET 100	
MG, 400 MG, 500 MG	.33
BRAFTOVI ORAL CAPSULE 50	
MG, 75 MG	34
BROVANA	.37
BRUKINSA	38
BUPHENYL ORAL TABLET	.39
buprenorphine HCl sublingual	.40
buprenorphine-naloxone sublingual	
film 12-3 mg, 2-0.5 mg, 4-1 mg, 8-2	
mg	40
BUSULFEX	40
CABOMETYX ORAL TABLET 20	. – 1
MG, 40 MG, 60 MG	42
calcitriol topical	
CALQUENCE	
capecitabine	
CAPLYTA	35
CAPRELSA ORAL TABLET 100	40
MG, 300 MG	
caspofungin	
CAYSTON	
cinacalcet	
clozapine oral tablet, disintegrating	.50
COMETRIQ ORAL CAPSULE 100	
MG/DAY(80 MG X1-20 MG X1),	
140 MG/DAY(80 MG X1-20 MG	
X3), 60 MG/DAY (20 MG X 3/DAY)	.51
COPAXONE SUBCUTANEOUS	
SYRINGE 20 MG/ML, 40 MG/ML	.52
SYRINGE 20 MG/ML, 40 MG/ML	53
SYRINGE 20 MG/ML, 40 MG/ML COPIKTRA	53 54
SYRINGE 20 MG/ML, 40 MG/ML COPIKTRA CORLANOR ORAL SOLUTION COSENTYX COSENTYX (2 SYRINGES)	53 54 55 55
SYRINGE 20 MG/ML, 40 MG/ML COPIKTRA CORLANOR ORAL SOLUTION COSENTYX COSENTYX (2 SYRINGES)	53 54 55 55
SYRINGE 20 MG/ML, 40 MG/ML COPIKTRA CORLANOR ORAL SOLUTION COSENTYX COSENTYX (2 SYRINGES) COSENTYX PEN	53 54 55 55 55
SYRINGE 20 MG/ML, 40 MG/ML COPIKTRA CORLANOR ORAL SOLUTION COSENTYX COSENTYX (2 SYRINGES) COSENTYX PEN COSENTYX PEN (2 PENS)	53 54 55 55 .55 .55
SYRINGE 20 MG/ML, 40 MG/ML COPIKTRA CORLANOR ORAL SOLUTION COSENTYX COSENTYX (2 SYRINGES) COSENTYX PEN COSENTYX PEN (2 PENS) COTELLIC	53 54 55 55 .55 .55 .55
SYRINGE 20 MG/ML, 40 MG/ML COPIKTRA CORLANOR ORAL SOLUTION COSENTYX COSENTYX (2 SYRINGES) COSENTYX PEN COSENTYX PEN (2 PENS) COTELLIC CYCLOSET	53 54 55 55 .55 .55 .55 .56 57
SYRINGE 20 MG/ML, 40 MG/ML COPIKTRA CORLANOR ORAL SOLUTION COSENTYX COSENTYX (2 SYRINGES) COSENTYX PEN COSENTYX PEN (2 PENS) COTELLIC CYCLOSET CYSTAGON	53 54 55 55 .55 .55 56 57 .58
SYRINGE 20 MG/ML, 40 MG/ML COPIKTRA CORLANOR ORAL SOLUTION COSENTYX COSENTYX (2 SYRINGES) COSENTYX PEN COSENTYX PEN (2 PENS) COTELLIC CYCLOSET CYSTAGON dalfampridine	53 54 55 55 .55 .55 .55 .56 57 .58 13
SYRINGE 20 MG/ML, 40 MG/ML COPIKTRA CORLANOR ORAL SOLUTION COSENTYX COSENTYX (2 SYRINGES) COSENTYX PEN COSENTYX PEN (2 PENS) COTELLIC CYCLOSET CYSTAGON dalfampridine DANYELZA	53 54 55 55 .55 .55 .55 .56 57 .58 13
SYRINGE 20 MG/ML, 40 MG/ML COPIKTRA CORLANOR ORAL SOLUTION COSENTYX COSENTYX (2 SYRINGES) COSENTYX PEN COSENTYX PEN (2 PENS) COTELLIC CYCLOSET CYSTAGON dalfampridine DANYELZA DAURISMO ORAL TABLET 100	53 54 55 55 55 55 55 57 57 58 13 59
SYRINGE 20 MG/ML, 40 MG/ML COPIKTRA CORLANOR ORAL SOLUTION COSENTYX. COSENTYX (2 SYRINGES) COSENTYX PEN. COSENTYX PEN (2 PENS). COTELLIC. CYCLOSET. CYSTAGON. dalfampridine. DANYELZA DAURISMO ORAL TABLET 100 MG, 25 MG.	53 54 55 55 .55 .55 57 .58 13 59 60
SYRINGE 20 MG/ML, 40 MG/ML COPIKTRA CORLANOR ORAL SOLUTION COSENTYX. COSENTYX (2 SYRINGES) COSENTYX PEN COSENTYX PEN (2 PENS) COTELLIC. CYCLOSET. CYSTAGON dalfampridine DANYELZA DAURISMO ORAL TABLET 100 MG, 25 MG deferiprone oral tablet 500 mg	53 54 55 55 .55 .55 57 .58 13 59 60
SYRINGE 20 MG/ML, 40 MG/ML COPIKTRA CORLANOR ORAL SOLUTION COSENTYX COSENTYX (2 SYRINGES) COSENTYX PEN COSENTYX PEN (2 PENS) COTELLIC CYCLOSET CYSTAGON dalfampridine DANYELZA DAURISMO ORAL TABLET 100 MG, 25 MG deferiprone oral tablet 500 mg DESCOVY ORAL TABLET 200-25	53 54 55 55 55 55 55 56 57 58 13 59 60 96
SYRINGE 20 MG/ML, 40 MG/ML COPIKTRA CORLANOR ORAL SOLUTION COSENTYX COSENTYX (2 SYRINGES) COSENTYX PEN COSENTYX PEN (2 PENS) COTELLIC CYCLOSET CYSTAGON dalfampridine DANYELZA DAURISMO ORAL TABLET 100 MG, 25 MG deferiprone oral tablet 500 mg DESCOVY ORAL TABLET 200-25 MG	53 54 55 55 55 55 55 55 57 57 58 13 59 60 96 61
SYRINGE 20 MG/ML, 40 MG/ML COPIKTRA CORLANOR ORAL SOLUTION COSENTYX COSENTYX (2 SYRINGES) COSENTYX PEN COSENTYX PEN (2 PENS) COTELLIC CYCLOSET CYSTAGON dalfampridine DANYELZA DAURISMO ORAL TABLET 100 MG, 25 MG deferiprone oral tablet 500 mg DESCOVY ORAL TABLET 200-25 MG DIACOMIT ORAL CAPSULE	53 54 55 55 55 55 55 57 57 59 60 96 61 .62
SYRINGE 20 MG/ML, 40 MG/ML COPIKTRA CORLANOR ORAL SOLUTION COSENTYX COSENTYX (2 SYRINGES) COSENTYX PEN COSENTYX PEN (2 PENS) COTELLIC CYCLOSET CYSTAGON dalfampridine DANYELZA DAURISMO ORAL TABLET 100 MG, 25 MG deferiprone oral tablet 500 mg DESCOVY ORAL TABLET 200-25 MG DIACOMIT ORAL CAPSULE DIAZEPAM INTENSOL	53 54 55 55 55 55 55 57 57 57 59 60 96 61 .62 30
SYRINGE 20 MG/ML, 40 MG/ML COPIKTRA CORLANOR ORAL SOLUTION COSENTYX COSENTYX (2 SYRINGES) COSENTYX PEN COSENTYX PEN (2 PENS) COTELLIC CYCLOSET CYSTAGON dalfampridine DANYELZA DAURISMO ORAL TABLET 100 MG, 25 MG deferiprone oral tablet 500 mg DESCOVY ORAL TABLET 200-25 MG DIACOMIT ORAL CAPSULE DIAZEPAM INTENSOL diazepam oral concentrate	53 54 55 55 55 55 55 55 57 .58 13 59 60 96 61 .62 .30 .30
SYRINGE 20 MG/ML, 40 MG/ML COPIKTRA CORLANOR ORAL SOLUTION COSENTYX COSENTYX (2 SYRINGES) COSENTYX PEN COSENTYX PEN (2 PENS) COTELLIC CYCLOSET CYSTAGON dalfampridine DANYELZA DAURISMO ORAL TABLET 100 MG, 25 MG deferiprone oral tablet 500 mg DESCOVY ORAL TABLET 200-25 MG DIACOMIT ORAL CAPSULE DIAZEPAM INTENSOL diazepam oral concentrate diclofenac sodium topical gel 3 %	53 54 55 55 55 55 55 55 57 .58 13 59 60 96 61 .62 .30 .30
SYRINGE 20 MG/ML, 40 MG/ML COPIKTRA CORLANOR ORAL SOLUTION COSENTYX COSENTYX (2 SYRINGES) COSENTYX PEN COSENTYX PEN (2 PENS) COTELLIC CYCLOSET CYSTAGON dalfampridine DANYELZA DAURISMO ORAL TABLET 100 MG, 25 MG deferiprone oral tablet 500 mg DESCOVY ORAL TABLET 200-25 MG DIACOMIT ORAL CAPSULE DIAZEPAM INTENSOL diazepam oral concentrate diclofenac sodium topical gel 3 % didanosine oral capsule,delayed	53 54 55 55 55 55 55 55 55 55 55 55 55 55
SYRINGE 20 MG/ML, 40 MG/ML COPIKTRA CORLANOR ORAL SOLUTION COSENTYX COSENTYX (2 SYRINGES) COSENTYX PEN COSENTYX PEN (2 PENS) COTELLIC CYCLOSET CYSTAGON dalfampridine DANYELZA DAURISMO ORAL TABLET 100 MG, 25 MG deferiprone oral tablet 500 mg DESCOVY ORAL TABLET 200-25 MG DIACOMIT ORAL CAPSULE DIAZEPAM INTENSOL diazepam oral concentrate diclofenac sodium topical gel 3 % didanosine oral capsule,delayed release(DR/EC) 250 mg, 400 mg	53 54 55 55 55 55 55 55 55 55 55 55 55 55
SYRINGE 20 MG/ML, 40 MG/ML COPIKTRA CORLANOR ORAL SOLUTION COSENTYX COSENTYX (2 SYRINGES) COSENTYX PEN COSENTYX PEN (2 PENS) COTELLIC CYCLOSET CYSTAGON dalfampridine DANYELZA DAURISMO ORAL TABLET 100 MG, 25 MG deferiprone oral tablet 500 mg DESCOVY ORAL TABLET 200-25 MG DIACOMIT ORAL CAPSULE DIAZEPAM INTENSOL diazepam oral concentrate diclofenac sodium topical gel 3 % didanosine oral capsule,delayed release(DR/EC) 250 mg, 400 mg DIFICID ORAL TABLET	53 54 55 55 55 55 55 55 55 55 55 55 55 55
SYRINGE 20 MG/ML, 40 MG/ML COPIKTRA CORLANOR ORAL SOLUTION COSENTYX COSENTYX (2 SYRINGES) COSENTYX PEN COSENTYX PEN (2 PENS) COTELLIC CYCLOSET CYSTAGON dalfampridine DANYELZA DAURISMO ORAL TABLET 100 MG, 25 MG deferiprone oral tablet 500 mg DESCOVY ORAL TABLET 200-25 MG DIACOMIT ORAL CAPSULE DIAZEPAM INTENSOL diazepam oral concentrate diclofenac sodium topical gel 3 % didanosine oral capsule,delayed release(DR/EC) 250 mg, 400 mg DIFICID ORAL TABLET dimethyl fumarate oral	53 54 55 55 55 55 55 55 55 55 55 55 55 55
SYRINGE 20 MG/ML, 40 MG/ML COPIKTRA CORLANOR ORAL SOLUTION COSENTYX COSENTYX (2 SYRINGES) COSENTYX PEN COSENTYX PEN (2 PENS) COTELLIC CYCLOSET CYSTAGON dalfampridine DANYELZA DAURISMO ORAL TABLET 100 MG, 25 MG deferiprone oral tablet 500 mg DESCOVY ORAL TABLET 200-25 MG DIACOMIT ORAL CAPSULE DIAZEPAM INTENSOL diazepam oral concentrate diclofenac sodium topical gel 3 % didanosine oral capsule, delayed release(DR/EC) 250 mg, 400 mg DIFICID ORAL TABLET dimethyl fumarate oral capsule, delayed release(DR/EC)	53 54 55 55 55 55 55 55 55 55 55 55 55 55
SYRINGE 20 MG/ML, 40 MG/ML COPIKTRA CORLANOR ORAL SOLUTION COSENTYX COSENTYX (2 SYRINGES) COSENTYX PEN COSENTYX PEN (2 PENS) COTELLIC CYCLOSET CYSTAGON dalfampridine DANYELZA DAURISMO ORAL TABLET 100 MG, 25 MG deferiprone oral tablet 500 mg DESCOVY ORAL TABLET 200-25 MG DIACOMIT ORAL CAPSULE DIAZEPAM INTENSOL diazepam oral concentrate diclofenac sodium topical gel 3 % didanosine oral capsule,delayed release(DR/EC) 250 mg, 400 mg DIFICID ORAL TABLET dimethyl fumarate oral capsule,delayed release(DR/EC) 120 mg, 120 mg (14)- 240 mg (46),	53 54 55 55 55 55 55 57 58 13 59 60 96 612 300 .30 .63 .645
SYRINGE 20 MG/ML, 40 MG/ML COPIKTRA CORLANOR ORAL SOLUTION COSENTYX COSENTYX (2 SYRINGES) COSENTYX PEN COSENTYX PEN (2 PENS) COTELLIC CYCLOSET CYSTAGON dalfampridine DANYELZA DAURISMO ORAL TABLET 100 MG, 25 MG deferiprone oral tablet 500 mg DESCOVY ORAL TABLET 200-25 MG DIACOMIT ORAL CAPSULE DIAZEPAM INTENSOL diazepam oral concentrate diclofenac sodium topical gel 3 % didanosine oral capsule, delayed release(DR/EC) 250 mg, 400 mg DIFICID ORAL TABLET dimethyl fumarate oral capsule, delayed release(DR/EC)	53 54 55 55 55 55 55 55 55 55 55 55 55 55

h	due a chine le vel con culo 10 mar 0 F	
bosentan	dronabinol oral capsule 10 mg, 2.5	~7
BOSULIF ORAL TABLET 100	mg, 5 mg	
<b>MG, 400 MG, 500 MG</b> 33	droxidopa	180
<b>BRAFTOVI ORAL CAPSULE 50</b>	DUPIXENT PEN	
<b>MG, 75 MG</b>	SUBCUTANEOUS PEN	
BROVANA	INJECTOR 300 MG/2 ML	.68
BRUKINSA	DUPIXENT SYRINGE	
BUPHENYL ORAL TABLET	SUBCUTANEOUS SYRINGE 200	
buprenorphine HCl sublingual40	MG/1.14 ML, 300 MG/2 ML	. 68
buprenorphine-naloxone sublingual	EDECRIN	
film 12-3 mg, 2-0.5 mg, 4-1 mg, 8-2	EDURANT	
<i>mg</i> 40	ELZONRIS	
BUSULFEX41	EMCYT	
CABOMETYX ORAL TABLET 20	EMEND ORAL CAPSULE 80 MG.	
MG, 40 MG, 60 MG42	EMEND ORAL CAPSULE,DOSE	.70
calcitriol topical	PACK	73
CALQUENCE 44	EMSAM	
capecitabine45		.75
CAPLYTA35 CAPRELSA ORAL TABLET 100	ENBREL SUBCUTANEOUS	70
	RECON SOLN	.76
<b>MG</b> , <b>300 MG</b>	ENBREL SUBCUTANEOUS	= 0
caspofungin47	SOLUTION	. 76
CAYSTON	ENBREL SUBCUTANEOUS	
cinacalcet49	SYRINGE 25 MG/0.5 ML (0.5), 50	
clozapine oral tablet, disintegrating50	MG/ML (1 ML)	
COMETRIQ ORAL CAPSULE 100	ENBREL SURECLICK	
MG/DAY(80 MG X1-20 MG X1),	ENDARI	
140 MG/DAY(80 MG X1-20 MG	ENSPRYNG	.78
X3), 60 MG/DAY (20 MG X 3/DAY).51	entecavir	. 25
COPAXONE SUBCUTANEOUS	EPCLUSA ORAL TABLET 400-	
SYRINGE 20 MG/ML, 40 MG/ML52	100 MG	. 79
<b>COPIKTRA</b>	EPIDIOLEX	.80
CORLANOR ORAL SOLUTION 54	EPIVIR HBV ORAL SOLUTION	
<b>COSENTYX</b>	EPZICOM	
COSENTYX (2 SYRINGES)	ERIVEDGE	
COSENTYX PEN	ERLEADA	
COSENTYX PEN (2 PENS)55	ERLOTINIB	
COTELLIC	ERTACZO	
<b>CYCLOSET</b>	ESBRIET	
CYSTAGON	EVENITY	
dalfampridine	everolimus (antineoplastic) oral	.07
<b>DANYELZA</b>	tablet 2.5 mg, 5 mg, 7.5 mg	7
DAURISMO ORAL TABLET 100		/
	everolimus (immunosuppressive)	
<b>MG</b> , <b>25 MG</b>	oral tablet 0.25 mg, 0.5 mg, 0.75	
deferiprone oral tablet 500 mg96	mg	
DESCOVY ORAL TABLET 200-25	EVOTAZ	
MG61	EVRYSDI	
DIACOMIT ORAL CAPSULE	EXJADE	
DIAZEPAM INTENSOL	FANAPT ORAL TABLET	. 92
diazepam oral concentrate30	FANAPT ORAL TABLETS, DOSE	
diclofenac sodium topical gel 3 %63	PACK	
didanosine oral capsule,delayed	FARYDAK	
release(DR/EC) 250 mg, 400 mg64	FASENRA PEN	. 94
DIFICID ORAL TABLET	fentanyl citrate buccal lozenge on a	
dimethyl fumarate oral	handle	. 95
capsule,delayed release(DR/EC)	FERRIPROX ORAL TABLET 500	
120 mg, 120 mg (14)- 240 mg (46),	MG	. 96
240 mg	FINTEPLA	
doxercalciferol oral	FIRDAPSE	
		-

fondaparinux subcutaneous syringe	Э
10 mg/0.8 mL, 2.5 mg/0.5 mL, 5 mg/0.4 mL, 7.5 mg/0.6 mL	100
FORTEO SUBCUTANEOUS PEN	100
INJECTOR 20 MCG/DOSE	
(600MCG/2.4ML)	101
fosamprenavir	
FOTIVDA	
FUZEON SUBCUTANEOUS	102
RECON SOLN	103
FYCOMPA ORAL SUSPENSION.	
FYCOMPA ORAL TABLET 10	104
MG, 12 MG, 2 MG, 4 MG, 6 MG, 8	
MG	104
GAMMAGARD LIQUID	
GAVRETO	
GENVOYA	
GILENYA	
GILOTRIF	
GLEOSTINE	
HARVONI ORAL TABLET 90-400	
MG	
HERCEPTIN HYLECTA	
HORIZANT	
HUMIRA PEN	
HUMIRA PEN CROHNS-UC-HS	
START	114
HUMIRA PEN PSOR-UVEITS-	
ADOL HS	114
HUMIRA SUBCUTANEOUS	
SYRINGE KIT 40 MG/0.8 ML	114
HUMIRA(CF)	
HUMIRA(CF) PEDI CROHNS	
STARTER SUBCUTANEOUS	
SYRINGE KIT 80 MG/0.8 ML, 80	
MG/0.8 ML-40 MG/0.4 ML	114
HUMIRA(CF) PEN	
HUMIRA(CF) PEN CROHNS-UC-	
HS	114
HUMIRA(CF) PEN PSOR-UV-	
ADOL HS	114
HYCAMTIN	
IBRANCE	
ICATIBANT	98
ICLUSIG ORAL TABLET 15 MG,	
45 MG	122
IDAMYCIN PFS	
IDHIFA	
imatinib oral tablet 100 mg, 400 mg	
	119
IMBRUVICA ORAL CAPSULE 140	
MG, 70 MG	120
IMBRUVICA ORAL TABLET	
INBRIJA	
INCRELEX	
INLYTA	
INQOVI	
INREBIC	126
INTELENCE ORAL TABLET 100	
MG, 200 MG, 25 MG	127

INTRON A INJECTION RECON	
SOLN	
INVIRASE ORAL TABLET	
ISENTRESS HD	132
ISENTRESS ORAL POWDER IN	
ISENTRESS ORAL TABLET	132
ISENTRESS ORAL	400
TABLET,CHEWABLE	
itraconazole oral capsule	133
JAKAFI ORAL TABLET 10 MG,	405
15 MG, 20 MG, 25 MG, 5 MG	135
JYNARQUE ORAL TABLETS,	
SEQUENTIAL 45 MG (AM)/ 15 MG	1
(PM), 60 MG (AM)/ 30 MG (PM),	400
90 MG (AM)/ 30 MG (PM)	
KALETRA	137
KALYDECO ORAL GRANULES	
IN PACKET 50 MG, 75 MG	
KALYDECO ORAL TABLET	
KISQALI	
KISQALI FEMARA CO-PACK	140
KUVAN ORAL	
TABLET,SOLUBLE	
lamivudine oral solution	142
lamivudine oral tablet 100 mg, 150	
mg, 300 mg	
lapatinib	281
LATUDA ORAL TABLET 120 MG,	
20 MG, 40 MG, 60 MG, 80 MG	
	146
LEUKINE INJECTION RECON	
SOLN	
leuprolide	
LEXIVA	149
lidocaine topical adhesive	4 5 0
patch,medicated 5 %	150
lidocaine topical ointment	
linezolid	
LONSURF	153
LORBRENA ORAL TABLET 100	. – .
MG, 25 MG	
LUPRON DEPOT (3 MONTH)	
LUPRON DEPOT (4 MONTH)	
LUPRON DEPOT (6 MONTH)	
LUPRON DEPOT-PED	
LUPRON DEPOT-PED (3 MONTH)	,
LYNPARZA MAKENA (PF)	
	4 5 0

# MAVENCLAD (10 TABLET PACK)

WAVENCLAD (10 TABLET PACK)	_
	9
MAVENCLAD (4 TABLET PACK) 15	9
MAVENCLAD (5 TABLET PACK) 15	9
<b>MAVENCLAD (6 TABLET PACK)</b> 15	9
MAVENCLAD (5 TABLET PACK) 15 MAVENCLAD (6 TABLET PACK) 15 MAVENCLAD (7 TABLET PACK) 15	9
MAVENCLAD (8 TABLET PACK) 15	9
MAVENCLAD (9 TABLET PACK) 15	a
MAVYRET ORAL TABLET	0
MAYZENT ORAL TABLET 0.25	0
MATZENT URAL TABLET 0.23	
<b>MG, 2 MG</b>	1
MAYZENT STARTER(FOR 2MG	
<b>MAINT)</b>	1
MEKINIST ORAL TABLET 0.5	
<b>MG, 2 MG</b>	2
<b>MEKTOVI</b> 16	3
<b>MENEST</b>	
miglustat16	
modafinil	6
MONJUVI	
MOVANTIK	
<b>MULPLETA</b>	
<b>NAGLAZYME</b> 17	
<b>NERLYNX</b> 17	
<b>NEULASTA</b>	
NEULASTA ONPRO 17	2
<b>NEUPOGEN</b>	3
nevirapine 174, 29	6
<b>NEXAVAR</b>	5
NEXLETOL	
NEXLIZET	
NINLARO	
NORPACE CR	
NORVIR ORAL SOLUTION	9
NOXAFIL	
NUBEQA	3
NUCALA SUBCUTANEOUS	
AUTO-INJECTOR 18	4
<b>NUCYNTA</b>	5
<b>NUCYNTA ER</b>	5
<b>NULOJIX</b>	6
NURTEC ODT	
octreotide acetate injection solution	-
	8
ODEFSEY	
<b>ODOMZO</b>	
OMNIPOD CLASSIC PDM	U
KIT(GEN 3)	1
OMNIPOD CLASSIC PODS (GEN	
<b>3)</b> 19	1
OMNIPOD DASH PDM KIT (GEN	
<b>4)</b> 19	1
OMNIPOD DASH PODS (GEN 4). 19	1
OMNITROPE	0
<b>ONUREG</b>	
ORFADIN ORAL CAPSULE 10	-
MG, 2 MG, 5 MG	3
ORIAHNN	
ORILISSA ORAL TABLET 150	-
	F
<b>MG, 200 MG</b> 19	C

ORKAMBI	.196
OTEZLA	
OTEZLA STARTER	197
oxandrolone oral tablet 10 mg, 2.5	;
mg	. 198
OXERVATE	. 199
OXLUMO	. 200
paliperidone oral tablet extended	
release 24hr 1.5 mg, 3 mg, 6 mg,	9
mg	. 129
PALYNZIQ	.201
pamidronate	. 202
PANRETIN	203
paricalcitol oral	. 204
penicillamine oral capsule	.205
, penicillamine oral tablet	
, PENTAM	
PERSERIS	
PIQRAY ORAL TABLET 200	
MG/DAY (200 MG X 1), 250	
MG/DAY (200 MG X1-50 MG X1),	
300 MG/DAY (150 MG X 2)	
PLEGRIDY SUBCUTANEOUS	
POMALYST	.210
POTELIGEO	
pramipexole oral tablet extended	
release 24 hr 0.375 mg, 0.75 mg,	
1.5 mg, 3 mg, 4.5 mg	.212
PREZCOBIX	
PREZISTA ORAL SUSPENSION	
PREZISTA ORAL TABLET 150	
MG, 600 MG, 75 MG, 800 MG	.214
<b>PROCRIT INJECTION SOLUTIOI</b>	N
10,000 UNIT/ML, 2,000 UNIT/ML,	
20,000 UNIT/ML, 3,000 UNIT/ML,	
4,000 UNIT/ML, 40,000 UNIT/ML.	.215
4,000 UNIT/ML, 40,000 UNIT/ML. PROMACTA ORAL POWDER IN	
4,000 UNIT/ML, 40,000 UNIT/ML. PROMACTA ORAL POWDER IN PACKET 12.5 MG	. 216
4,000 UNIT/ML, 40,000 UNIT/ML. PROMACTA ORAL POWDER IN PACKET 12.5 MG PROMACTA ORAL TABLET	. 216 .216
4,000 UNIT/ML, 40,000 UNIT/ML. PROMACTA ORAL POWDER IN PACKET 12.5 MG PROMACTA ORAL TABLET PULMOZYME	. 216 .216
4,000 UNIT/ML, 40,000 UNIT/ML. PROMACTA ORAL POWDER IN PACKET 12.5 MG PROMACTA ORAL TABLET PULMOZYME quetiapine oral tablet extended	. 216 .216 .218
4,000 UNIT/ML, 40,000 UNIT/ML. PROMACTA ORAL POWDER IN PACKET 12.5 MG PROMACTA ORAL TABLET PULMOZYME quetiapine oral tablet extended release 24 hr	. 216 .216 .218 .238
4,000 UNIT/ML, 40,000 UNIT/ML. PROMACTA ORAL POWDER IN PACKET 12.5 MG PROMACTA ORAL TABLET PULMOZYME quetiapine oral tablet extended release 24 hr raloxifene	. 216 .216 .218 .238 36
4,000 UNIT/ML, 40,000 UNIT/ML. PROMACTA ORAL POWDER IN PACKET 12.5 MG PROMACTA ORAL TABLET PULMOZYME quetiapine oral tablet extended release 24 hr raloxifene REGRANEX.	. 216 .216 .218 . 238 36 219
4,000 UNIT/ML, 40,000 UNIT/ML. PROMACTA ORAL POWDER IN PACKET 12.5 MG PROMACTA ORAL TABLET PULMOZYME quetiapine oral tablet extended release 24 hr raloxifene REGRANEX. RELENZA DISKHALER.	. 216 .216 .218 . 238 36 219 .220
4,000 UNIT/ML, 40,000 UNIT/ML. PROMACTA ORAL POWDER IN PACKET 12.5 MG PROMACTA ORAL TABLET PULMOZYME quetiapine oral tablet extended release 24 hr raloxifene REGRANEX RELENZA DISKHALER REMODULIN.	. 216 .216 .218 . 238 36 219 .220 . 221
4,000 UNIT/ML, 40,000 UNIT/ML. PROMACTA ORAL POWDER IN PACKET 12.5 MG PROMACTA ORAL TABLET PULMOZYME. quetiapine oral tablet extended release 24 hr raloxifene. REGRANEX. RELENZA DISKHALER. REMODULIN REPATHA PUSHTRONEX.	. 216 .218 . 238 36 219 . 220 . 221 222
4,000 UNIT/ML, 40,000 UNIT/ML. PROMACTA ORAL POWDER IN PACKET 12.5 MG PROMACTA ORAL TABLET PULMOZYME quetiapine oral tablet extended release 24 hr raloxifene REGRANEX RELENZA DISKHALER REMODULIN REPATHA PUSHTRONEX. REPATHA SURECLICK	. 216 .218 . 218 36 219 . 220 . 221 222 222
4,000 UNIT/ML, 40,000 UNIT/ML. PROMACTA ORAL POWDER IN PACKET 12.5 MG PROMACTA ORAL TABLET PULMOZYME quetiapine oral tablet extended release 24 hr raloxifene REGRANEX RELENZA DISKHALER REMODULIN REPATHA PUSHTRONEX REPATHA SURECLICK REPATHA SYRINGE	. 216 .218 . 238 36 219 .220 .221 222 .222 . 222
4,000 UNIT/ML, 40,000 UNIT/ML. PROMACTA ORAL POWDER IN PACKET 12.5 MG PROMACTA ORAL TABLET PULMOZYME quetiapine oral tablet extended release 24 hr raloxifene REGRANEX. RELENZA DISKHALER REMODULIN REPATHA PUSHTRONEX. REPATHA SURECLICK REPATHA SYRINGE. RETROVIR INTRAVENOUS	. 216 .218 . 238 36 .219 .220 . 221 .222 .222 . 222 . 223
4,000 UNIT/ML, 40,000 UNIT/ML. PROMACTA ORAL POWDER IN PACKET 12.5 MG PROMACTA ORAL TABLET PULMOZYME quetiapine oral tablet extended release 24 hr raloxifene REGRANEX. RELENZA DISKHALER REMODULIN. REPATHA PUSHTRONEX. REPATHA SURECLICK REPATHA SYRINGE. RETROVIR INTRAVENOUS REVLIMID.	. 216 .218 . 238 36 .219 .220 . 221 .222 .222 . 222 . 223
4,000 UNIT/ML, 40,000 UNIT/ML. PROMACTA ORAL POWDER IN PACKET 12.5 MG PROMACTA ORAL TABLET PULMOZYME quetiapine oral tablet extended release 24 hr raloxifene REGRANEX RELENZA DISKHALER REMODULIN REPATHA PUSHTRONEX REPATHA SURECLICK REPATHA SYRINGE RETROVIR INTRAVENOUS REVLIMID REYATAZ ORAL CAPSULE 200	. 216 .218 . 238 36 219 .220 .221 .222 .222 .222 .223 .223
4,000 UNIT/ML, 40,000 UNIT/ML. PROMACTA ORAL POWDER IN PACKET 12.5 MG PROMACTA ORAL TABLET PULMOZYME quetiapine oral tablet extended release 24 hr raloxifene REGRANEX REGRANEX RELENZA DISKHALER REMODULIN. REPATHA PUSHTRONEX REPATHA SURECLICK. REPATHA SYRINGE RETROVIR INTRAVENOUS REVLIMID. REYATAZ ORAL CAPSULE 200 MG, 300 MG	. 216 .218 . 238 36 219 .220 .221 .222 .222 .222 .223 .223
4,000 UNIT/ML, 40,000 UNIT/ML. PROMACTA ORAL POWDER IN PACKET 12.5 MG	. 216 .218 . 238 36 219 . 220 . 221 . 222 . 222 . 222 . 223 . 224 . 225
4,000 UNIT/ML, 40,000 UNIT/ML. PROMACTA ORAL POWDER IN PACKET 12.5 MG PROMACTA ORAL TABLET PULMOZYME quetiapine oral tablet extended release 24 hr raloxifene REGRANEX. RELENZA DISKHALER REMODULIN REPATHA PUSHTRONEX. REPATHA SURECLICK REPATHA SURECLICK REPATHA SYRINGE. RETROVIR INTRAVENOUS REVLIMID REYATAZ ORAL CAPSULE 200 MG, 300 MG REYATAZ ORAL POWDER IN PACKET	. 216 .218 . 238 36 219 .220 .221 .222 .222 .222 .223 .224 . 225 . 225
4,000 UNIT/ML, 40,000 UNIT/ML. PROMACTA ORAL POWDER IN PACKET 12.5 MG PROMACTA ORAL TABLET PULMOZYME quetiapine oral tablet extended release 24 hr raloxifene REGRANEX. RELENZA DISKHALER REMODULIN. REPATHA PUSHTRONEX. REPATHA SURECLICK REPATHA SYRINGE RETROVIR INTRAVENOUS REVLIMID REYATAZ ORAL CAPSULE 200 MG, 300 MG REYATAZ ORAL POWDER IN PACKET RIABNI	. 216 .218 . 238 36 219 .220 .221 .222 .222 .222 .223 .224 . 225 .225 .225
4,000 UNIT/ML, 40,000 UNIT/ML. PROMACTA ORAL POWDER IN PACKET 12.5 MG PROMACTA ORAL TABLET PULMOZYME quetiapine oral tablet extended release 24 hr raloxifene REGRANEX RELENZA DISKHALER REMODULIN REPATHA PUSHTRONEX REPATHA SURECLICK REPATHA SYRINGE RETROVIR INTRAVENOUS REVLIMID REVLIMID REYATAZ ORAL CAPSULE 200 MG, 300 MG REYATAZ ORAL POWDER IN PACKET RIABNI ribavirin oral capsule	. 216 .218 . 238 36 219 .220 .221 .222 .222 .222 .223 .224 . 225 .225 .225 .225 .226 .227
4,000 UNIT/ML, 40,000 UNIT/ML. PROMACTA ORAL POWDER IN PACKET 12.5 MG PROMACTA ORAL TABLET PULMOZYME quetiapine oral tablet extended release 24 hr raloxifene REGRANEX RELENZA DISKHALER REMODULIN REPATHA PUSHTRONEX REPATHA SURECLICK REPATHA SYRINGE RETROVIR INTRAVENOUS REVLIMID REYATAZ ORAL CAPSULE 200 MG, 300 MG REYATAZ ORAL POWDER IN PACKET RIABNI ribavirin oral capsule ribavirin oral tablet 200 mg	. 216 .218 .238 36 219 .220 .221 .222 .222 .222 .223 .224 .225 .225 .225 .225 .225 .225 .227 .227
4,000 UNIT/ML, 40,000 UNIT/ML. PROMACTA ORAL POWDER IN PACKET 12.5 MG PROMACTA ORAL TABLET PULMOZYME quetiapine oral tablet extended release 24 hr raloxifene REGRANEX RELENZA DISKHALER REMODULIN REPATHA PUSHTRONEX REPATHA SURECLICK REPATHA SYRINGE RETROVIR INTRAVENOUS REVLIMID REVLIMID REYATAZ ORAL CAPSULE 200 MG, 300 MG REYATAZ ORAL POWDER IN PACKET RIABNI ribavirin oral capsule	. 216 .218 .238 36 219 .220 .221 .222 .222 .222 .223 .224 .225 .225 .225 .225 .225 .225 .227 .227

ROZLYTREK ORAL CAPSULE	
100 MG, 200 MG	230
RUBRACA	231
rufinamide oral suspension	24
RYDAPT	232
RYDAPT SABRIL ORAL TABLET	233
SAMSCA	234
SANCUSO	
SANDOSTATIN LAR DEPOT	
INTRAMUSCULAR	
SUSPENSION, EXTENDED REL	
RECON	235
SAPHRIS SUBLINGUAL TABLE	:
10 MG, 2.5 MG, 5 MG	
sapropterin oral tablet, soluble	
SECUADO	
SELZENTRY ORAL TABLET 15	n 200
MG, 25 MG, 300 MG, 75 MG	
SEROQUEL XR ORAL TABLET	237
EXTENDED RELEASE 24 HR	000
SIGNIFOR LAR	
sildenafil (pulm.hypertension) ora	
tablet	240
SKYRIZI SUBCUTANEOUS	~
SYRINGE 75 MG/0.83 ML	241
SKYRIZI SUBCUTANEOUS	
sodium phenylbutyrate oral tablet	
SOFOSBUVIR-VELPATASVIR	
SOMATULINE DEPOT	
Somavert subcutaneous recon so	oln
10 mg	
SOMAVERT SUBCUTANEOUS	244
SOMAVERT SUBCUTANEOUS RECON SOLN 15 MG, 20 MG, 2	244 <b>5</b>
SOMAVERT SUBCUTANEOUS	244 <b>5</b>
SOMAVERT SUBCUTANEOUS RECON SOLN 15 MG, 20 MG, 2	244 <b>5</b> 244
SOMÁVERT SUBCUTANEOUS RECON SOLN 15 MG, 20 MG, 2 MG, 30 MG	244 5 244 245
SOMÁVERT SUBCUTANEOUS RECON SOLN 15 MG, 20 MG, 2 MG, 30 MG SPRYCEL	244 5 244 245 246
SOMÁVERT SUBCUTANEOUS RECON SOLN 15 MG, 20 MG, 2 MG, 30 MG SPRYCEL stavudine oral capsule STIVARGA	244 5 244 245 246 247
SOMAVERT SUBCUTANEOUS RECON SOLN 15 MG, 20 MG, 2 MG, 30 MG SPRYCEL stavudine oral capsule STIVARGA STRIBILD	244 5 244 245 246 247 248
SOMAVERT SUBCUTANEOUS RECON SOLN 15 MG, 20 MG, 2 MG, 30 MG SPRYCEL stavudine oral capsule STIVARGA STRIBILD sunitinib	244 5 244 245 246 247 248 251
SOMAVERT SUBCUTANEOUS RECON SOLN 15 MG, 20 MG, 2 MG, 30 MG SPRYCEL stavudine oral capsule STIVARGA STRIBILD sunitinib SUNOSI	244 5 244 245 246 247 248 248 251 249
SOMÁVERT SUBCUTANEOUS RECON SOLN 15 MG, 20 MG, 2 MG, 30 MG SPRYCEL stavudine oral capsule STIVARGA STRIBILD sunitinib SUNOSI SUSTIVA	244 5 244 245 246 247 248 251 249 250
SOMÁVERT SUBCUTANEOUS RECON SOLN 15 MG, 20 MG, 2 MG, 30 MG SPRYCEL stavudine oral capsule STIVARGA STRIBILD sunitinib SUNOSI SUSTIVA SUTENT	244 5 244 245 246 247 248 251 249 250 251
SOMÁVERT SUBCUTANEOUS RECON SOLN 15 MG, 20 MG, 2 MG, 30 MG SPRYCEL stavudine oral capsule STIVARGA STRIBILD sunitinib SUNOSI SUSTIVA SUTENT SYNERA	244 5 244 245 246 247 248 251 250 251 252
SOMAVERT SUBCUTANEOUS RECON SOLN 15 MG, 20 MG, 2 MG, 30 MG SPRYCEL stavudine oral capsule STIVARGA STRIBILD sunitinib SUNOSI SUSTIVA SUTENT SYNERA TABLOID	244 5 244 245 246 247 248 251 249 250 251 252 253
SOMAVERT SUBCUTANEOUS RECON SOLN 15 MG, 20 MG, 2 MG, 30 MG SPRYCEL stavudine oral capsule STIVARGA STRIBILD sunitinib SUNOSI SUSTIVA SUTENT SYNERA TABLOID tacrolimus topical	244 5 244 245 246 247 248 251 250 251 252 253 217
SOMAVERT SUBCUTANEOUS RECON SOLN 15 MG, 20 MG, 2 MG, 30 MG SPRYCEL stavudine oral capsule STIVARGA STRIBILD SUNOSI SUSTIVA SUTENT SYNERA TABLOID tacrolimus topical tadalafil (pulm. hypertension)	244 5 244 245 246 247 248 251 250 251 252 252 253 217 4
SOMAVERT SUBCUTANEOUS RECON SOLN 15 MG, 20 MG, 2 MG, 30 MG SPRYCEL stavudine oral capsule STIVARGA STRIBILD SUNOSI SUSTIVA SUTENT SYNERA TABLOID tacrolimus topical tadalafil (pulm. hypertension) TAFINLAR	244 5 244 245 246 247 248 251 250 251 252 252 253 217 4
SOMÁVERT SUBCUTANEOUS RECON SOLN 15 MG, 20 MG, 2 MG, 30 MG SPRYCEL stavudine oral capsule STIVARGA STRIBILD sunitinib SUNOSI SUSTIVA SUTENT SYNERA TABLOID tacrolimus topical tadalafil (pulm. hypertension) TAFINLAR TAGRISSO ORAL TABLET 40	244 <b>5</b> 244 245 246 247 248 251 250 251 252 253 217 4 254
SOMÁVERT SUBCUTANEOUS RECON SOLN 15 MG, 20 MG, 2 MG, 30 MG SPRYCEL stavudine oral capsule STIVARGA STRIBILD sunitinib SUNOSI SUSTIVA SUTENT SYNERA TABLOID tacrolimus topical tadalafil (pulm. hypertension) TAFINLAR TAGRISSO ORAL TABLET 40 MG, 80 MG	244 <b>5</b> 244 245 246 247 248 251 250 251 252 253 217 4 254
SOMÁVERT SUBCUTANEOUS RECON SOLN 15 MG, 20 MG, 2 MG, 30 MG SPRYCEL stavudine oral capsule STIVARGA STRIBILD sunitinib SUNOSI SUSTIVA SUTENT SYNERA TABLOID tacrolimus topical tadalafil (pulm. hypertension) TAFINLAR TAGRISSO ORAL TABLET 40 MG, 80 MG TAKHZYRO SUBCUTANEOUS	244 <b>5</b> 244 245 246 247 248 251 251 250 253 217 4 254 255
SOMAVERT SUBCUTANEOUS RECON SOLN 15 MG, 20 MG, 2 MG, 30 MG SPRYCEL stavudine oral capsule STIVARGA STRIBILD sunitinib SUNOSI SUSTIVA SUTENT SYNERA TABLOID tacrolimus topical tadalafil (pulm. hypertension) TAFINLAR TAGRISSO ORAL TABLET 40 MG, 80 MG TAKHZYRO SUBCUTANEOUS SOLUTION	244 5 244 245 246 247 248 251 250 251 252 253 217 4 254 255 255 256
SOMÁVERT SUBCUTANEOUS RECON SOLN 15 MG, 20 MG, 2 MG, 30 MG SPRYCEL stavudine oral capsule STIVARGA STRIBILD SUNOSI SUSTIVA SUTENT SYNERA TABLOID tacrolimus topical tadalafil (pulm. hypertension) TAFINLAR TAGRISSO ORAL TABLET 40 MG, 80 MG TAKHZYRO SUBCUTANEOUS SOLUTION TALZENNA ORAL CAPSULE 0.	244 5 244 245 247 248 251 249 250 251 252 253 254 255 256 25
SOMÁVERT SUBCUTANEOUS RECON SOLN 15 MG, 20 MG, 2 MG, 30 MG SPRYCEL stavudine oral capsule STIVARGA STRIBILD SUNOSI SUNOSI SUSTIVA SUTENT SYNERA TABLOID tacrolimus topical tadalafil (pulm. hypertension) TAFINLAR TAGRISSO ORAL TABLET 40 MG, 80 MG TAKHZYRO SUBCUTANEOUS SOLUTION TALZENNA ORAL CAPSULE 0. MG, 1 MG	244 5 244 245 246 247 248 251 250 251 252 253 254 255 256 25 257
SOMAVERT SUBCUTANEOUS RECON SOLN 15 MG, 20 MG, 2 MG, 30 MG SPRYCEL stavudine oral capsule STIVARGA STRIBILD sunitinib SUNOSI SUNOSI SUSTIVA SUTENT SYNERA TABLOID tacrolimus topical tadalafil (pulm. hypertension) TAFINLAR TAGRISSO ORAL TABLET 40 MG, 80 MG TAKHZYRO SUBCUTANEOUS SOLUTION TALZENNA ORAL CAPSULE 0. MG, 1 MG	244 <b>5</b> 244 245 246 247 248 251 250 251 252 253 255 256 <b>25</b> 257 36
SOMÁVERT SUBCUTANEOUS RECON SOLN 15 MG, 20 MG, 2 MG, 30 MG SPRYCEL stavudine oral capsule STIVARGA STRIBILD sunitinib SUNOSI SUNOSI SUSTIVA SUTENT SYNERA TABLOID tacrolimus topical tadalafil (pulm. hypertension) TAFINLAR TAGRISSO ORAL TABLET 40 MG, 80 MG TAKHZYRO SUBCUTANEOUS SOLUTION TALZENNA ORAL CAPSULE 0. MG, 1 MG tamoxifen TARCEVA	244 <b>5</b> 244 245 246 247 248 251 250 251 252 253 254 255 256 <b>25</b> 257 36 258
SOMÁVERT SUBCUTANEOUS RECON SOLN 15 MG, 20 MG, 2 MG, 30 MG SPRYCEL stavudine oral capsule STIVARGA STRIBILD sunitinib SUNOSI SUSTIVA SUTENT SYNERA TABLOID tacrolimus topical tadalafil (pulm. hypertension) TAFINLAR TAGRISSO ORAL TABLET 40 MG, 80 MG TAKHZYRO SUBCUTANEOUS SOLUTION TALZENNA ORAL CAPSULE 0. MG, 1 MG tamoxifen TARCEVA TARGRETIN TOPICAL	244 <b>5</b> 244 245 246 247 248 251 250 251 252 253 254 255 256 <b>25</b> 257 36 258
SOMÁVERT SUBCUTANEOUS RECON SOLN 15 MG, 20 MG, 2 MG, 30 MG SPRYCEL stavudine oral capsule STIVARGA STRIBILD sunitinib SUNOSI SUSTIVA SUTENT SYNERA TABLOID tacrolimus topical tadalafil (pulm. hypertension) TAFINLAR TAGRISSO ORAL TABLET 40 MG, 80 MG TAKHZYRO SUBCUTANEOUS SOLUTION TALZENNA ORAL CAPSULE 0. MG, 1 MG tamoxifen TARGRETIN TOPICAL TARGRETIN TOPICAL TASIGNA ORAL CAPSULE 150	244 5 244 245 247 248 251 250 251 252 253 254 255 256 25 256 25 36 258 259
SOMAVERT SUBCUTANEOUS RECON SOLN 15 MG, 20 MG, 2 MG, 30 MG SPRYCEL stavudine oral capsule STIVARGA STRIBILD sunitinib SUNOSI SUSTIVA SUTENT SYNERA TABLOID tacrolimus topical tadalafil (pulm. hypertension) TAFINLAR TAGRISSO ORAL TABLET 40 MG, 80 MG TAKHZYRO SUBCUTANEOUS SOLUTION	244 5 244 245 246 247 248 251 250 250 251 252 253 255 256 25 256 25 258 258 259 260

TAZVERIK	
temozolomide	
tenofovir disoproxil fumarate	
ТЕРМЕТКО	
THALOMID	
THYMOGLOBULIN	
TIBSOVO	.268
TIGAN INTRAMUSCULAR	269
TIVICAY	.270
tobramycin in 0.225 % NaCl	271
TOPOŠAR	272
TRACLEER	
tranexamic acid intravenous	274
tretinoin (antineoplastic)	.275
TRIKAFTA ORAL TABLETS,	
SEQUENTIAL 100-50-75 MG(D)	
150 MG (N)	276
TRINTELLIX	
TRIUMEQ	
TRUVADA	
TURALIO	
TYKERB	
TYMLOS	
UBRELVY	
VALCHLOR	
VALCYTE ORAL RECON SOLN.	284
valganciclovir oral tablet	.284
vancomycin oral capsule	.285
VARUBI	88
VENCLEXTA	
VENCLEXTA STARTING PACK	.287
VENTAVIS	
VERQUVO	
VERZENIO	
VIBERZI	
vigabatrin oral powder in packet	233
VIIBRYD ORAL TABLET	
VIIBRYD ORAL TABLETS,DOSE	
PACK 10 MG (7)- 20 MG (23)	292
VILTEPSO	
VIMPAT ORAL SOLUTION	
VIMPAT ORAL TABLET	294
VIMPAT ORAL TABLETS, DOSE	004
PACK VIRACEPT ORAL TABLET	.294
VIREAD VITRAKVI ORAL CAPSULE	291
VIZIMPRO	
voriconazole oral	
VOTRIENT	
VUMERITY	
VYVANSE ORAL CAPSULE	
VYXEOS	
WAKIX	
XALKORI	
XCOPRI	308
XCOPRI MAINTENANCE PACK	200
ORAL TABLET 250MG/DAY(150	
MG X1-100MG X1), 350 MG/DAY	
(200 MG X1-150MG X1)	308

XCOPRI TITRATION PACK	200
XELJANZ ORAL SOLUTION	
XELJANZ ORAL TABLET	
XELJANZ XR	
XERMELO	88
XIFAXAN ORAL TABLET 200	
MG, 550 MG	. 310
XOLAIR	. 311
XOSPATA	312
XPOVIO ORAL TABLET 60MG	
TWICE WEEK (120 MG/WEEK),	
80MG TWICE WEEK (160	
MG/WEEK)	. 313
XTANDI ORAL CAPSULE	
XYREM	
YONSA	
ZAVESCA	
ZEJULA	
ZELBORAF	
ZEMAIRA	
ZIAGEN ORAL SOLUTION	321
zidovudine	322
zoledronic acid-mannitol-water	
intravenous piggyback 5 mg/100	
mL	. 323
ZOLINZA	. 324
<b>ZORTRESS ORAL TABLET 0.25</b>	-
MG, 0.5 MG, 0.75 MG	325
ZYDELIG	
ZYFLO	
ZYKADIA ORAL TABLET	
ZYTIGA ORAL TABLET 250 MG	
500 MG	,
vvv m v	. 520

This formulary was updated on 01/01/2022. For more recent information or other questions, please contact Health First Health Plans Customer Service at 1.800.716.7737 or, for TTY users, 1.800.955.8771, weekdays from 8 a.m. to 8 p.m. and Saturdays from 8 a.m. to noon. From October 1 through March 31, we are available seven days a week from 8 a.m. to 8 p.m. or visit myHFHP.org.

Customer Service has language interpreter services available for non-English speakers at no cost.

This information is also available at no cost in other formats. By contacting Customer Service you may request your materials be read aloud, emailed, or mailed in large print.

You must generally use network pharmacies to use your prescription drug benefit. Benefits, formulary, pharmacy network, premium and/or copayments/coinsurance may change on January 1, 2021, and from time to time during the year.

Health First Health Plans is an HMO plan with a Medicare contract. Enrollment in Health First Health Plans depends on contract renewal.

The Formulary, pharmacy network, may change at any time. You will receive notice when necessary.



6450 U.S. Highway 1 Rockledge, FL 32955 myHFHP.org

# **Nondiscrimination Notice**

Health First Health Plans complies with applicable Federal civil rights laws and does not discriminate on the basis of race, color, national origin, age, disability or sex. Health First Health Plans does not exclude people or treat them differently because of race, color, national origin, age, disability or sex.

Health First Health Plans:

- Provides free aids and services to people with disabilities to communicate effectively with us, such as:
  - Qualified sign language interpreters
  - Written information in other formats (large print, accessible electronic formats)
- Provides free language services to people whose primary language is not English, such as:
  - Qualified interpreters
  - Information written in other languages

If you need these services, please contact our Civil Rights Coordinator.

If you believe that Health First Health Plans has failed to provide these services or discriminated in another way on the basis of race, color, national origin, age, disability, or sex, you can file a grievance with: Civil Rights Coordinator, 6450 US Highway 1, Rockledge, FL 32955, 321-434-4521, 1-800-955-8771 (TTY), Fax: 321-434-4362, <u>civilrightscoordinator@HF.org</u>. You can file a grievance in person or by mail, fax or email. If you need help filing a grievance our Civil Rights Coordinator, is available to help you.

You can also file a civil rights complaint with the U.S. Department of Health and Human Services, Office for Civil Rights electronically through the Office for Civil Rights Complaint Portal, available at <u>https://ocrportal.hhs.gov/ocr/portal/lobby.jsf</u>, or by mail or phone at: U.S. Department of Health and Human Services, 200 Independence Avenue SW., Room 509F, HHH Building, Washington, DC 20201, 1-800-368-1019, 800-537-7697 (TDD).

Complaint forms are available at <u>http://www.hhs.gov/ocr/office/file/index.html</u>.

Health First Health Plans is an HMO plan with a Medicare contract. Enrollment in Health First Health Plans depends on contract renewal.

Y0089\_MPINFO7271\_C (04/19)



6450 U.S. Highway 1 Rockledge, FL 32955 myHFHP.org

**English:** ATTENTION: If you speak English, language assistance services, free of charge, are available to you. Call 1-800-716-7737 (TTY: 1-800-955-8771).

**Spanish:** ATENCIÓN: si habla español, tiene a su disposición servicios gratuitos de asistencia lingüística. Llame al 1-800-716-7737 (TTY: 1-800-955-8771).

**French Creole:** ATANSYON: Si w pale Kreyòl Ayisyen, gen sèvis èd pou lang ki disponib gratis pou ou. Rele 1-800-716-7737 (TTY: 1-800-955-8771).

Vietnamese: CHÚ Ý: Nếu bạn nói Tiếng Việt, có các dịch vụ hỗ trợ ngôn ngữ miễn phí dành cho bạn. Gọi số 1-800-716-7737 (TTY: 1-800-955-8771).

**Portuguese:** ATENÇÃO: Se fala português, encontram-se disponíveis serviços linguísticos, grátis. Ligue para 1-800-716-7737 (TTY: 1-800-955-8771).

Chinese: 注意:如果您使用繁體中文,您可以免費獲得語言援助服務。請致電 1-800-716-7737(TTY: 1-800-955-8771)。

**French:** ATTENTION : Si vous parlez français, des services d'aide linguistique vous sont proposés gratuitement. Appelez le 1-800-716-7737 (ATS : 1-800-955-8771).

**Tagalog:** PAUNAWA: Kung nagsasalita ka ng Tagalog, maaari kang gumamit ng mga serbisyo ng tulong sa wika nang walang bayad. Tumawag sa 1-800-716-7737 (TTY: 1-800-955-8771).

**Russian:** ВНИМАНИЕ: Если вы говорите на русском языке, то вам доступны бесплатные услуги перевода. Звоните 1-800-716-7737 (телетайп: 1-800-955-8771).

# Arabic:

ملحوظة: إذا كنت تتحدث اذكر اللغة، فإن خدمات المساعدة اللغوية تتوافر لك بالمجان. اتصل برقم 1-7737-716-800 (رقم هاتف الصم والبكم: 1-877-850-800).

**Italian:** ATTENZIONE: In caso la lingua parlata sia l'italiano, sono disponibili servizi di assistenza linguistica gratuiti. Chiamare il numero 1-800-716-7737 (TTY: 1-800-955-8771).

**German:** ACHTUNG: Wenn Sie Deutsch sprechen, stehen Ihnen kostenlos sprachliche Hilfsdienstleistungen zur Verfügung. Rufnummer: 1-800-716-7737 (TTY: 1-800-955-8771).

Korean: 주의: 한국어를 사용하시는 경우, 언어 지원 서비스를 무료로 이용하실 수 있습니다. 1-800-716-7737 (TTY: 1-800-955-8771)번으로 전화해 주십시오.

**Polish:** UWAGA: Jeżeli mówisz po polsku, możesz skorzystać z bezpłatnej pomocy językowej. Zadzwoń pod numer 1-800-716-7737 (TTY: 1-800-955-8771).

Gujarati: સુચના: જો તમે ગુજરાતી બોલતા હો, તો નિ:શુલ્ક ભાષા સહાય સેવાઓ તમારા માટે ઉપલબ્ધ છે. ફોન કરો 1-800-716-7737 (TTY: 1-800-955-8771).

Thai: เรียน: ถ้าคุณพูดภาษาไทยคุณสามารถใช้บริการช่วยเหลือทางภาษาได้ฟรี โทร 1-800-716-7737 (TTY: 1-800-955-8771).

Health First Health Plans is an HMO plan with a Medicare contract. Enrollment in Health First Health Plans depends on contract renewal. Y0089\_MPINFO6466 (08/17)